

A Regularization Corrected Score Method for Nonlinear Regression Models with Covariate Error

David M. Zucker^{1,*}, Malka Gorfine², Yi Li³, Mahlet G. Tadesse⁴, and Donna Spiegelman⁵

¹Department of Statistics, Hebrew University, Mount Scopus, Jerusalem, Israel

²Faculty of Industrial Engineering and Management, Technion - Israel Institute of Technology, Haifa, Israel

³Department of Biostatistics, University of Michigan, Ann Arbor MI, USA

⁴Department of Mathematics and Statistics, Georgetown University, Washington DC, USA

⁵Departments of Epidemiology and Biostatistics, Harvard School of Public Health, Boston MA, USA

**email*: mszucker@mscc.huji.ac.il

SUMMARY: Many regression analyses involve explanatory variables that are measured with error, and failing to account for this error is well known to lead to biased point and interval estimates of the regression coefficients. We present here a new general method for adjusting for covariate error. Our method consists of an approximate version of the Stefanski-Nakamura corrected score approach, using the method of regularization to obtain an approximate solution of the relevant integral equation. We develop the theory in the setting of classical likelihood models; this setting covers, for example, linear regression, nonlinear regression, logistic regression, and Poisson regression. The method is extremely general in terms of the types of measurement error models covered, and is a functional method in the sense of not involving assumptions on the distribution of the true covariate. We discuss the theoretical properties of the method and present simulation results in the logistic regression setting (univariate and multivariate). For illustration, we apply the method to data from the Harvard Nurses' Health Study concerning the relationship between physical activity and breast cancer mortality in the period following a diagnosis of breast cancer.

KEY WORDS: Errors in variables; nonlinear models; logistic regression; integral equations

This paper has been submitted for consideration for publication in *Biometrics*

1. Introduction

Many regression analyses involve explanatory variables that are measured with error. Failing to account for the covariate error leads to biased estimates of the regression coefficients, and hence a large literature has developed on the problem of correcting for covariate measurement error. Fuller (1987) provides an authoritative account for linear models, while Carroll et al. (2006) summarizes the literature on nonlinear models. The nonlinear case remains an active research area, bearing on such common statistical models as nonlinear regression with a continuous response, logistic regression for binary responses, Poisson regression for count data, and Cox regression for survival data. This paper presents a flexible new method for nonlinear regression problems with covariate error, built on earlier work.

Three basic study designs are of interest: (1) the replicate measures design, where repeat covariate measurements are available (either for all individuals or for a subsample), (2) the internal validation design, where the true covariate values are available on a sample of individuals in the main study, and (3) the external validation design, where the key parameters of the measurement error distribution are estimated (assuming reasonable transportability) from an external study, independent of the main study, with paired measurements of the true and surrogate covariate. Also, two types of methods are of interest: structural methods, which involve model assumptions on the distribution of the true covariates, and functional methods, which do not require such assumptions.

Various approaches have been proposed. We focus on the SIMEX and corrected score approaches, which are functional modeling approaches. These are general approaches that can handle both internal and external validation designs as well as, with slight adaptation, the replicate measures design. Our proposed method is based on the corrected score approach.

The SIMEX method of Cook and Stefanski (1994) involves simulating new covariate values with various levels of artificially added measurement error, carrying out a naive model fit for

each of the resulting new data sets, and then back-extrapolating to zero measurement error. While some success has been achieved with this approach, obviously the back-extrapolation process is uncertain. Moreover, SIMEX can be hard to apply in certain non-classical settings. One challenging setting is when the measurement error variance depends on the true covariate value. A version of SIMEX that can handle this setting under the replicate measures design was developed by Devanarayan and Stefanski (2002), but it does not appear that SIMEX can handle this setting under internal or external validation study designs.

The corrected score approach, advanced by Stefanski (1989) and Nakamura (1990, 1992), involves replacing the likelihood score in the conventional likelihood-based analysis with a function of the surrogate covariates that serves as an unbiased substitute. In Section 2, we present the exact definition. For the independent additive error model, Nakamura (1990) showed that this approach yields consistent estimators in the case of normal linear regression, Poisson regression, Gamma regression, and inverse Gaussian regression. Nakamura (1992) presented an approximate corrected score method for the Cox regression model, which Kong and Gu (1999) later showed to yield consistent estimates. Novick and Stefanski (2002) presented a corrected score method that is aimed at the independent additive error model with normal errors, and is valid when the likelihood score function is an entire function in the complex plane (such as, for example, Poisson regression with an exponential link). Stefanski, Novick, and Devanarayan (2005) elaborate on this approach. When the error-prone covariate is discrete, a corrected score can be formulated easily; the relevant theory was developed for generalized linear models by Akazawa, Kinukawa, and Nakamura (1998), and extended to the Cox regression model by Zucker and Spiegelman (2008).

On the other hand, for logistic regression with additive normal error, Stefanski (1989) showed that an exact corrected score method does not exist. Huang and Wang (2001) presented an exact modified corrected score method for logistic regression, with clever

reweighting of the terms in the original score function so that an exact corrected score can be found. This reweighting leads to some loss in efficiency. Moreover, the method of Huang and Wang is designed only for the case of independent additive measurement error. Buzas (2009) presents an approximate corrected score method for the logistic regression model with high efficiency when the covariate effect is moderate, but this method is designed only for the case of independent additive normal error.

The basic problem with the corrected score approach in the logistic regression model and other cases with a continuous error-prone covariate X is that obtaining the corrected score requires solving a challenging integral equation. The equation involved is a Fredholm integral equation of the first kind, discussed by Delves and Mohamed (1985, Ch. 12) and Kress (1989, Ch. 16). Such equations do not always have an exact solution; the logistic regression problem is one case of this. Moreover, even when an exact solution exists, the problem can be ill-conditioned. We tried to tackle the case of a continuous covariate by discretizing the covariate to various degrees of fineness and applying the methodology for the discrete case, but with only limited success. The classification matrix tended to be ill-conditioned even with a modest degree of fineness, such as six categories.

In this paper, we develop a new approach. The idea is to handle the integral equation using the method of regularization (Delves and Mohamed, 1985, Sec. 12.3; Kress, 1989, Ch. 16), which involves minimizing a penalized distance function to obtain an approximate solution. In contrast with the original integral equation problem, the regularized problem always has a solution, and is reasonably well conditioned provided that the weight α on the penalty term is not too small. As α tends to infinity, the estimation procedure tends to a naive analysis in which we ignore the covariate error, and simply substitute the surrogate covariate value for the true value. Conversely, under suitable conditions, as α tends to zero the procedure approaches an exact corrected score procedure. The idea is to push α as close as possible to

zero to get good estimates of the model parameters. We call our approach the regularized corrected score (RECS) approach.

The advantage of RECS is that it is extremely flexible. Its formulation is very general, and it is a functional method with no modeling of the distribution of the true covariate, but only of the conditional density of the surrogate covariate given the true covariate. The method can handle both internal and external validation designs. It can handle the replicate measures design as well, with the overall surrogate measurement defined as the sample mean (or other summary measure) of the available measurements on the individual. Moreover, the method can handle arbitrary measurement error structures, not just independent additive measurement error. Differential measurement error, where the measurement error depends on the response, is also covered. The goal of this paper is to develop the RECS method in detail for the classical likelihood setting. Section 2 lays out the setting and background. Section 3 presents the proposed procedure and its theoretical properties. Section 4 presents simulation results under the logistic regression model. Section 5 presents a real-data illustration of the method in the logistic regression setting. Section 6 presents a brief discussion.

2. Setting and Background

We assume a typical setup with n independent units whose response values Y_i , $i = 1, \dots, n$, follow a regression model involving several covariates. We assume for now that only one of the covariates is subject to error; later we will generalize to the case of multiple error-prone covariates. We denote by X_i the true value of the error-prone covariate, and by W_i the measured value. We let \mathbf{Z}_i denote the vector of error-free covariates, which may include an arbitrary number of discrete and continuous components. We denote the conditional density or mass function of Y_i given (X_i, \mathbf{Z}_i) by $f(y|X_i, \mathbf{Z}_i, \boldsymbol{\theta})$, where $\boldsymbol{\theta}$ is a p -vector of unknown parameters, including regression coefficients and auxiliary parameters such as error variances. In contrast with the SIMEX method, the Huang and Wang (2001) method, and most other

methods in the literature, we provide the option of allowing the measurement error to depend on X_i , \mathbf{Z}_i , and the outcome Y_i (differential error). We denote by $a_i(x, w)$ the conditional density of W_i given $X_i = x$, with the subscript i signifying possible dependence on \mathbf{Z}_i and Y_i . To ease the presentation of the theoretical results, we assume that (X_i, \mathbf{Z}_i) are i.i.d. random vectors. We stress, however, that our method does not involve any modeling (either parametric or nonparametric) of the distribution of (X_i, \mathbf{Z}_i) . The theory can, in principle, be extended to the case where (X_i, \mathbf{Z}_i) are non-random values satisfying suitable ergodicity conditions.

The classical normalized likelihood score function when there is no covariate error is given by $\mathbf{U}(\boldsymbol{\theta}) = n^{-1} \sum_i \mathbf{u}_i(X_i, \boldsymbol{\theta})$, where $\mathbf{u}_i(x, \boldsymbol{\theta}) = \mathbf{u}(Y_i, x, \mathbf{Z}_i, \boldsymbol{\theta})$ with $\mathbf{u}(y, x, \mathbf{z}, \boldsymbol{\theta}) = [\partial/\partial\boldsymbol{\theta}] \log f(y|x, \mathbf{z}, \boldsymbol{\theta})$. The maximum likelihood estimate (MLE) is obtained by solving the equation $\mathbf{U}(\boldsymbol{\theta}) = \mathbf{0}$. The idea of the Stefanski-Nakamura corrected score approach is to find a function $\bar{\mathbf{u}}(y, w, \mathbf{z}, \boldsymbol{\theta})$ such that

$$E[\bar{\mathbf{u}}(Y_i, W_i, \mathbf{Z}_i, \boldsymbol{\theta}) | X_i, \mathbf{Z}_i, Y_i] = \mathbf{u}(Y_i, X_i, \mathbf{Z}_i, \boldsymbol{\theta}). \quad (1)$$

We define $\bar{\mathbf{u}}_i(w, \boldsymbol{\theta}) = \bar{\mathbf{u}}(Y_i, w, \mathbf{Z}_i, \boldsymbol{\theta})$, and then use the modified likelihood score function $\bar{\mathbf{U}}(\boldsymbol{\theta}) = n^{-1} \sum_i \bar{\mathbf{u}}_i(W_i, \boldsymbol{\theta})$ in place of $\mathbf{U}(\boldsymbol{\theta})$ as the basis for estimation. The estimation equation thus becomes $\bar{\mathbf{U}}(\boldsymbol{\theta}) = \mathbf{0}$.

In the present setting, equation (1) for the corrected score function takes the form

$$\int a_i(x, w) \bar{u}_{ij}(w) dw = u_{ij}(x), \quad (2)$$

where $u_{ij}(x)$ and $\bar{u}_{ij}(w)$, respectively, denote the j -th component of $\mathbf{u}_i(x, \boldsymbol{\theta})$ and $\bar{\mathbf{u}}_i(w, \boldsymbol{\theta})$ (suppressing the argument $\boldsymbol{\theta}$ in the definitions), and the integral is over the entire range of W . As indicated in the introduction, we do not seek an exact solution to (2), but instead use the method of regularization to find an approximate solution.

Define the integral operator $A_i g(x) = \int a_i(x, w) g(w) dw$. Write $\Delta_{ij}(x) = u_{ij}(x) - A_i u_{ij}(x)$

and $\bar{\Delta}_{ij}(x) = \bar{u}_{ij}(x) - u_{ij}(x)$. We can then write (2) as $A_i \bar{\Delta}_{ij} = \Delta_{ij}$. We seek the $\bar{\Delta}_{ij}(\cdot, \boldsymbol{\theta}, \alpha)$ that minimizes the penalized loss function

$$\mathcal{L}_{ij}(\bar{\Delta}_{ij}) = \|A_i \bar{\Delta}_{ij} - \Delta_{ij}\|^2 + \alpha \|\bar{\Delta}_{ij}\|^2, \quad (3)$$

where $\|g\|^2$ denotes the weighted squared L^2 norm $\|g\|^2 = \int c(v)g(v)^2 dv$ and $\alpha > 0$ is a penalty factor. The penalty is introduced in order to avoid ill-conditioning and ensure existence of a solution. After obtaining $\bar{\Delta}_{ij}(\cdot, \boldsymbol{\theta}, \alpha)$, we use $\bar{u}_{ij}(x, \boldsymbol{\theta}, \alpha) = u_{ij}(x) + \bar{\Delta}_{ij}(x, \boldsymbol{\theta}, \alpha)$ as a corrected score term.

For the weight function $c(v)$, we propose as a generic choice the standard normal density, i.e., $c(v) = \varphi(v)$ with $\varphi(v) = \exp(-v^2/2)/\sqrt{2\pi}$, after standardizing W to mean 0 and variance 1. The weight function is designed to put emphasis on the region of the covariate space where the bulk of the data lie. One could consider tailoring the choice of the weight function to the pattern of the observed distribution of W , but we do not discuss this here.

We formulate the minimization problem in terms of $\bar{\Delta}_{ij}(\cdot, \boldsymbol{\theta}, \alpha)$ rather than $\bar{u}_{ij}(\cdot, \boldsymbol{\theta}, \alpha)$ in order to anchor the procedure at $u_{ij}(w)$, which corresponds to the naive analysis where we ignore the covariate error and substitute W_i for X_i . See Hansen (1994, Sec. 2, second paragraph) for the idea of centering the regularization process around an initial estimate of the desired solution to the integral equation. As $\alpha \rightarrow \infty$, the loss function $\mathcal{L}_{ij}(\bar{\Delta}_{ij})$ puts increasingly heavy weight on $\|\bar{\Delta}_{ij}\|^2$, causing the minimizer $\bar{\Delta}_{ij}$ to tend to 0 and thus leading to the naive estimates based on $u_{ij}(w)$. At the other extreme, as $\alpha \rightarrow 0$, the problem of minimizing $\mathcal{L}_{ij}(\bar{\Delta}_{ij})$ approaches the problem of solving $A_i \bar{\Delta}_{ij} = \Delta_{ij}$. We first describe the procedure for a fixed α , and then discuss the selection of the value of α .

By using the L^2 norm, we ensure that the problem of minimizing the loss function always has a unique solution, and the solution has a convenient form. Delves and Mohamed (1985, Sec. 12.3) and Kress (1989, Theorem 16.1) present the relevant theory. Let $A_i^* h(w) = \int a_i(x, w)h(x)dx$ denote the operator adjoint of A_i . Then, for any L^2 function δ , the minimizer

of $\mathcal{L}(\bar{\delta}; A_i, \delta, \alpha) = \|A_i\bar{\delta} - \delta\| + \alpha\|\bar{\delta}\|^2$ is given by $\bar{\delta} = (A_i^*A_i + \alpha\mathcal{I})^{-1}A_i^*\delta$, where \mathcal{I} is the identity operator. The next section describes a numerical scheme for finding the solution that leads to a simple linear system of equations. We thus obtain an easily implemented procedure.

3. The Procedure

3.1 The Procedure for a Given α

To numerically determine the minimizer of $\mathcal{L}_{ij}(\bar{\Delta}_{ij})$, we use a Galerkin-type basis expansion approach, in the spirit of (though not identical to) Delves and Mohamed (1985, Sec. 12.4).

Specifically, we represent the solution $\bar{\Delta}_{ij}(\cdot, \boldsymbol{\theta}, \alpha)$ in a basis expansion

$$\bar{\Delta}_{ij}(x, \boldsymbol{\theta}, \alpha) = \sum_{m=1}^M d_{ijm}(\alpha)\psi_m(x), \quad (4)$$

where the ψ_m are specified basis functions. In our numerical work, we use the ‘‘probabilists’’ Hermite polynomials, which are orthonormal with respect to the weight function φ . One has to choose the number M of basis functions to include. We found that $M = 6$ yields good performance; the results with $M = 4$ are inferior to those with $M = 6$, while the results with $M = 10$ are similar to those with $M = 6$ but with more outliers.

Denote $\phi_{im}(x) = A_i\psi_m(x)$ and $\mathbf{d}_{ij} = [d_{ij1} \dots d_{ijM}]^T$ (suppressing the argument α in d_{ijm} for the time being). We then can express the objective function $\mathcal{L}_{ij}(\bar{\Delta}_{ij})$ as

$$\mathcal{L}_{ij}(\bar{\Delta}_{ij}) = \left\| \sum_{m=1}^M d_{ijm}\phi_{im} - \Delta_{ij} \right\|^2 + \alpha \mathbf{d}_{ij}^T \mathbf{d}_{ij},$$

where $\|\bar{\Delta}_{ij}\|^2 = \mathbf{d}_{ij}^T \mathbf{d}_{ij}$ because of the orthonormality of the ψ_m functions. We now approximate the L^2 norm in the first term on the right side via the quadrature approximation

$$\int \varphi(v)g(v)dv \doteq \sum_{k=1}^K q_k g(x_k), \quad (5)$$

where x_k and q_k are the classical Gauss-Hermite quadrature points and weights (modified slightly to account for our use of the weight function $\exp(-v^2/2)/\sqrt{2\pi}$ as opposed to the standard Hermite weight function $\exp(-v^2)$). Given the approximation (5), we can express

the objective function as

$$\begin{aligned}\mathcal{L}_{ij}(\bar{\Delta}_{ij}) &= \sum_{k=1}^K q_k \left[\sum_{m=1}^M d_{ijm} \phi_{im}(x_k) - \Delta_{ij}(x_k) \right]^2 + \alpha \mathbf{d}_{ij}^T \mathbf{d}_{ij} \\ &= \sum_{k=1}^K \left[\sum_{m=1}^M d_{ijm} \tilde{\phi}_{imk} - \tilde{\Delta}_{ijk} \right]^2 + \alpha \mathbf{d}_{ij}^T \mathbf{d}_{ij},\end{aligned}$$

where $\tilde{\phi}_{imk} = \sqrt{q_k} \phi_m(x_k)$ and $\tilde{\Delta}_{ijk} = \sqrt{q_k} \Delta_{ij}(x_k)$. Next, define the matrix $\tilde{\Phi}_i$ by $\tilde{\Phi}_{ikm} = \tilde{\phi}_{imk}$ and the vector $\tilde{\Delta}_{ij} = [\tilde{\Delta}_{ij1} \dots \tilde{\Delta}_{ijK}]$. We obtain $\mathcal{L}_{ij}(\bar{\Delta}_{ij}) = (\tilde{\Phi}_i \mathbf{d}_{ij} - \tilde{\Delta}_{ij})^T (\tilde{\Phi}_i \mathbf{d}_{ij} - \tilde{\Delta}_{ij}) + \alpha \mathbf{d}_{ij}^T \mathbf{d}_{ij}$. We then find, by standard least squares theory, that the vector $\mathbf{d}_{ij}(\alpha)$ that minimizes the above quantity is given by $\mathbf{d}_{ij}(\alpha) = \mathbf{C}(\alpha) \tilde{\Delta}_{ij}$, where $\mathbf{C}(\alpha) = (\tilde{\Phi}_i^T \tilde{\Phi}_i + \alpha \mathbf{I})^{-1}$. Note that $\mathbf{C}(\alpha)$ does not depend on $\boldsymbol{\theta}$. Finally, we define $\bar{u}_{ij}(w, \boldsymbol{\theta}, \alpha) = u_{ij}(w, \boldsymbol{\theta}) + \bar{\Delta}_{ij}(w, \boldsymbol{\theta}, \alpha)$, where $\bar{\Delta}_{ij}(w, \boldsymbol{\theta}, \alpha)$ is given by (4) with $d_{ijm}(\alpha)$ as just described. Then, as indicated in the preceding section, we put $\bar{\mathbf{U}}(\boldsymbol{\theta}, \alpha) = n^{-1} \sum_i \bar{\mathbf{u}}_i(W_i, \boldsymbol{\theta}, \alpha)$ and define the estimator $\hat{\boldsymbol{\theta}}^{(\alpha)}$ to be the solution to $\bar{\mathbf{U}}(\boldsymbol{\theta}, \alpha) = \mathbf{0}$.

In the course of the foregoing procedure, we have to evaluate integrals of the form

$$E[g(W_i) | X_i = x] = A_i g(x) = \int a_i(x, w) g(w) dw. \quad (6)$$

Integrals of this type appear in $\phi_{im}(x) = A_i \psi_m(x)$ and in $\Delta_{ij}(x) = u_{ij}(x) - A_i u_{ij}(x)$. These integrals can be evaluated by K' -point numerical quadrature for suitable K' . Web Appendix A.1 presents the details. In regard to the choice of K and K' , in our numerical work we generally used $K = K' = 20$; we reran selected simulations with $K = K' = 30$ and obtained similar results. In a data analysis, the analyst can try a succession of increasing values of K and K' , and stop when there is no further change in the results.

In practice, $a_i(x, w)$ has to be estimated, using data from an internal or external validation study (or a replicate measures study). We assume that $a_i(x, w)$ follows a known parametric model depending on parameters $\boldsymbol{\xi}$ (distinct from $\boldsymbol{\theta}$) which are estimated from the relevant data. We thus write $a_i(x, w, \boldsymbol{\xi})$. The parametric model is allowed to be of any specified form.

Thus, in addition to the classical independent additive error model, we allow models with dependence between the error and the true covariate value, and models with differential error. We have examined the effect of misspecifying the parametric form. In practice, the parametric model would be chosen based on analysis of the validation data. Regression methods can be used to model $E[W|X]$. Heterogeneity of variance can be assessed and modeled by applying regression methods to the squares of the residuals, as in Davidian and Carroll (1987). These models can be checked using standard tools. The distributional form for the error can be selected by examining the standardized residuals from the regression of W on X .

3.2 Theoretical Properties

In general, $\hat{\boldsymbol{\theta}}^{(\alpha)}$ will not converge to the true value $\boldsymbol{\theta}_0$ of $\boldsymbol{\theta}$, but rather to a limit $\tilde{\boldsymbol{\theta}}^{(\alpha)}$ that is close to $\boldsymbol{\theta}_0$ when α is small. We cannot make α arbitrarily small, but we can try to make it small enough to obtain estimates with small bias, and the numerical studies in the next section indicate that this goal can be achieved. Thus, our method does not produce an exactly consistent estimator, but it does produce an approximately consistent estimator. Moreover, under standard regularity conditions, $\sqrt{n}(\hat{\boldsymbol{\theta}}^{(\alpha)} - \tilde{\boldsymbol{\theta}}^{(\alpha)})$ is asymptotically normal. These properties are formalized in the following theorem, proved in Web Appendix A.2.

Theorem: Define $\bar{\mathbf{u}}_E(\boldsymbol{\theta}, \alpha) = E[\bar{\mathbf{u}}_i(W_i, \boldsymbol{\theta}, \alpha)]$. Let $\bar{\mathbf{D}}^{(\alpha)}(\boldsymbol{\theta})$ be the Jacobian of $-\bar{\mathbf{U}}(\boldsymbol{\theta}, \alpha)$. Then, under the regularity conditions A1-A4 in Web Appendix A.2, the following results hold.

a. We have $E[\bar{u}_{ij}(W_i, \boldsymbol{\theta}, \alpha)|X_i, \mathbf{Z}_i, Y_i] = u_{ij}(X_i, \boldsymbol{\theta}) + r_{ij}(X_i, \boldsymbol{\theta}, \alpha)$ with $r_{ij}(\cdot, \boldsymbol{\theta}, \alpha)$ converging uniformly in $\boldsymbol{\theta}$ to zero as $\alpha \rightarrow 0$.

b. Similarly, with $u_{ijs}(x, \boldsymbol{\theta}) = [\partial/\partial\theta_s]u_{ij}(x, \boldsymbol{\theta})$ and $\bar{u}_{ijs}(w, \boldsymbol{\theta}, \alpha) = [\partial/\partial\theta_s]\bar{u}_{ij}(w, \boldsymbol{\theta}, \alpha)$, we have $E[\bar{u}_{ijs}(W_i, \boldsymbol{\theta}, \alpha)|X_i, \mathbf{Z}_i, Y_i] = u_{ijs}(X_i, \boldsymbol{\theta}) + r_{ijs}(X_i, \boldsymbol{\theta}, \alpha)$ with $r_{ijs}(\cdot, \boldsymbol{\theta}, \alpha)$ converging uniformly in $\boldsymbol{\theta}$ to zero as $\alpha \rightarrow 0$.

c. For all α sufficiently small, the equation $\bar{\mathbf{u}}_E(\boldsymbol{\theta}, \alpha) = \mathbf{0}$ has a unique solution, which we denote by $\tilde{\boldsymbol{\theta}}^{(\alpha)}$. For fixed α , we have $\hat{\boldsymbol{\theta}}^{(\alpha)} \rightarrow \tilde{\boldsymbol{\theta}}^{(\alpha)}$ almost surely as $n \rightarrow \infty$.

d. We have $\tilde{\boldsymbol{\theta}}^{(\alpha)} \rightarrow \boldsymbol{\theta}_0$ as $\alpha \rightarrow 0$.

e. If $a_i(x, w)$ is known, $\sqrt{n}(\hat{\boldsymbol{\theta}}^{(\alpha)} - \tilde{\boldsymbol{\theta}}^{(\alpha)})$ is asymptotically mean-zero normal with covariance matrix that can be estimated using the sandwich estimator

$$\mathbf{V}^{(\alpha)}(\hat{\boldsymbol{\theta}}^{(\alpha)}) = \bar{\mathbf{D}}^{(\alpha)}(\hat{\boldsymbol{\theta}}^{(\alpha)})^{-1} \mathbf{F}^{(\alpha)}(\hat{\boldsymbol{\theta}}^{(\alpha)}) \bar{\mathbf{D}}^{(\alpha)}(\hat{\boldsymbol{\theta}}^{(\alpha)})^{-1}$$

with $\mathbf{F}^{(\alpha)}(\boldsymbol{\theta}) = n^{-1} \sum_{i=1}^n \bar{\mathbf{u}}_i(\boldsymbol{\theta}, \alpha) \bar{\mathbf{u}}_i(\boldsymbol{\theta}, \alpha)^T$. Under a parametric model for $a_i(x, w)$ with estimated parameters $\boldsymbol{\xi}$, a similar result holds, with a suitable adjustment to the estimated covariance matrix to account for the estimation of $\boldsymbol{\xi}$, as described in Web Appendix A.3.

Remark: Assumptions A1-A3 are typical assumptions made in asymptotic theory; see, for example, van der Vaart (1998, p. 46, bottom). Assumption A4 is a modest assumption that holds in many cases of interest. For example, suppose (X, W) follows the independent additive error model $W = X + \sigma\varepsilon$, where ε is a random variable with density f_ε , independent of X . We then have $a(x, w) = \sigma^{-1} f_\varepsilon((w - x)/\sigma) = \sigma^{-1} \tilde{f}_\varepsilon((x - w)/\sigma)$, with $\tilde{f}_\varepsilon(u) = f_\varepsilon(-u)$. The assumption thus will be satisfied provided that the location-scale family of densities $f(x; w, \sigma) = \sigma^{-1} \tilde{f}_\varepsilon((x - w)/\sigma)$ is a complete family of densities with respect to the parameters (w, σ) . This condition certainly holds if f_ε is a density of exponential family form; see Lehmann (1986, p. 142). Next, consider the extended model $W = X + \sigma(X, \boldsymbol{\gamma})\varepsilon$, which we examine in our numerical studies, where $\boldsymbol{\gamma}$ is a vector of parameters. Assumption A4 will hold in this setting if the family of densities

$$f(x; w, \boldsymbol{\gamma}) = \frac{1}{\sigma(x, \boldsymbol{\gamma})} \tilde{f}_\varepsilon\left(\frac{x - w}{\sigma(x, \boldsymbol{\gamma})}\right) \Big/ \int \frac{1}{\sigma(x', \boldsymbol{\gamma})} \tilde{f}_\varepsilon\left(\frac{x' - w}{\sigma(x', \boldsymbol{\gamma})}\right) dx'$$

is a complete family of densities with respect to the parameters $(w, \boldsymbol{\gamma})$. Again, this condition will hold if f_ε is a density of exponential family form.

3.3 Choice of the Penalty Parameter α

The issue of how to choose the penalty parameter in a regularization problem has been investigated in previous literature. Hansen (1994, 2007) describes three leading criteria: the L-curve criterion, the GCV criterion, and the quasi-optimality criterion. We tried all three, and found the GCV criterion to be the most satisfactory. The GCV criterion is

$$\text{GCV}(\alpha) = \frac{(\tilde{\Phi}_i \mathbf{d}_{ij}(\alpha) - \tilde{\Delta}_{ij})^T (\tilde{\Phi}_i \mathbf{d}_{ij}(\alpha) - \tilde{\Delta}_{ij})}{[\text{trace}(\mathbf{I} - \mathbf{C}(\alpha))]^2},$$

and α is chosen to minimize this quantity. In our setting, with a separate value $\text{GCV}_{ij}(\alpha)$ for each i and j , we use the summary criterion $\text{GCV}^*(\alpha) = (np)^{-1} \sum_{i=1}^n \sum_{j=1}^p \text{GCV}_{ij}(\alpha)$. In implementing this rule, we evaluate $\tilde{\Delta}_{ij}$ at the naive estimate of $\boldsymbol{\theta}$, and then keep α fixed at the resulting value for the remainder of the estimation process.

3.4 Multiple Error-Prone Covariates

The method can be extended to the case of two error-prone covariates X_1 and X_2 . For the basis functions, we use the tensor product of the univariate basis functions. The integrals in quantities of the form $A_i g(x)$ become double integrals, which are evaluated by bivariate quadrature. In the L^2 norm appearing in the objective function, we take the weight function to be $c(w_1, w_2) = \varphi(w_1)\varphi(w_2)$, and evaluate the integral using the bivariate version of (5).

In the case of three or more error-prone covariates, the situation becomes more complicated. Taking the basis function set to be the tensor product of the univariate basis functions will typically produce too large a basis function set, so some reduction will be necessary. One option is to take the basis function set to include all the univariate basis functions for the individual covariates plus the cross-products of the linear terms. In the evaluation of the integrals $A_i g(x) = E[g(\mathbf{W}_i) | X_i = x]$, a Monte-Carlo method will probably be more workable than a classical quadrature method. The L^2 norms can be computed using a multivariate version of (5), but the computational load may be demanding.

The simulation work presented in the next section includes results for the case of two error-prone covariates. On a practical level, it appears challenging to apply our method with three or more error-prone covariates. However, many applications involve only one or two error-prone covariates, and thus can be handled by our method in a reasonable way. An arbitrary number of error-free covariates can be handled without difficulty.

4. The Logistic Regression Model: Simulation Studies

4.1 Simulation Study Designs

To investigate the finite sample performance of our method, we conducted a series of simulation studies. This subsection describes the simulation study designs; the next subsection describes the results. The studies were conducted in the setting of the logistic regression model. The response variable Y_i equals either 0 or 1. Defining $\mathbf{T}_i = (\mathbf{X}_i, \mathbf{Z}_i)$ and $T_{i0} \equiv 1$, the model is logit $\Pr(Y_i = 1 | \mathbf{X}_i = \mathbf{x}, \mathbf{Z} = \mathbf{z}) = \sum_{j=1}^p \beta_{j-1} t_{j-1}$ and the score function $\mathbf{u}(y, x, z, \boldsymbol{\theta})$ is

$$u_j(y, x, z, \boldsymbol{\theta}) = t_{j-1} \left[y - \text{expit} \left(\sum_{\ell=1}^p \beta_{\ell-1} t_{\ell-1} \right) \right], \quad \text{expit}(a) = e^a / (1 + e^a).$$

We examined the following methods: (1) naive analysis ignoring measurement error, (2) RECS, (3) the Novick and Stefanski (2001) complex variable corrected score method (N&S), (4) SIMEX, (5) the Huang and Wang (2001) nonparametric corrected score method (H&W), (6) a Bayesian MCMC-type method (at the suggestion of a referee), patterned after Richardson et al. (2002). The SIMEX method was applied with linear (SIMEX-L), quadratic (SIMEX-Q), and nonlinear extrapolation (SIMEX-NL). We show the results for SIMEX-Q in the tables, and the results for the other versions in the web appendix. For the Huang and Wang method, which requires replicate measurements of W , we took two replicates per individual and doubled the error variance for comparability with the other methods. Note that H&W is designed to provide accurate estimates only for the slope, not for the intercept. All simulation

results for Methods 1-5 are based on 1,000 simulation replications. The MCMC method is very time-consuming, so for this method we ran only 100 replications.

The SIMEX method is designed for independent additive normal measurement error. The N&S method is designed for the case where the measurement error is independent additive normal and the likelihood score function is an entire function in the complex plane. The latter condition does not hold for logistic regression, and thus the N&S method is not designed to handle any of the simulation scenarios we have studied. The H&W nonparametric method is designed for independent additive measurement error with an arbitrary distribution, which is not modeled in any way. The RECS method is designed for parametric measurement error models of arbitrary form, including non-normal, heteroscedastic, and differential error.

The details of the Bayesian MCMC method are presented in Web Appendix B. The MCMC algorithm was run for 100,000 iterations, with the last 40,000 iterations used for inference. Our adaptation of the Richardson et al. method involves an input parameter η that regulates the variance of the prior distributions on β_0 and β_1 . Specifically, the prior on β_j was taken to be normal with mean equal to the naive estimate and variance equal to η times the estimated variance of the naive estimate. We present results for $\eta = 100$. Results for $\eta = 10$ are provided in the web appendix. It was necessary for the prior on β_0 and β_1 to be moderately informative to ensure convergence of the MCMC sampler. This is an inherent difficulty with the Bayesian approach in the context of the main study / external validation study design that we consider, with no individuals for which the outcome Y and the true covariate X are both observed. The setting $\eta = 100$ corresponds to the maximum degree of noninformativeness under which convergence of the MCMC sampler is maintained across all scenarios studied.

For each estimator, we summarize the bias in terms of the mean and median of the difference between the estimated and true parameter values, and the dispersion in terms of the empirical standard deviation and standardized interquartile range (dividing the raw

interquartile range by 1.349, which is the ratio between the interquartile range and the standard deviation for a normal distribution). The median and the interquartile range are more robust to outliers, and thus provide further insight in addition to that provided by the mean and the standard deviation. In addition, we present the coverage rates of the 95% Wald confidence interval based on asymptotic normal theory. For the naive method, variance estimates were computed using the standard asymptotic variance formula. For the RECS and N&S methods, an explicit sandwich-type formula was used. For SIMEX, we used the jackknife estimator in the R package `simex`, while for H&W we used a bootstrap variance estimator. For the Bayesian MCMC method, interval estimation was based on 95% highest posterior density (HPD) intervals (Box and Tiao, 1973, Section 2.8).

In Simulation Sets A and B, we considered the case of a single continuous error-prone covariate X_i and no other covariates. In these two simulation sets, we worked in the setting of a main study / external validation study design, involving a main study sample with data on W and Y and an external validation sample with data on W and X . The main study sample size was 200 and the external validation sample size was 70. The measurement error parameters were estimated by maximum likelihood. The RECS method was implemented with $M = 6$ basis functions and $K = K' = 20$ quadrature points. The true values of the regression parameters were set at $\beta_0 = \beta_1 = 1$.

Simulation Set A involved measurement error models of the form $W_i = X_i + \epsilon_i$, where ϵ_i is normally distributed, but with error variance possibly depending on X_i and Y_i . We examined three simulation scenarios, as follows:

Scenario A1: $X_i \sim N(0, 1)$, $\epsilon_i | (X_i, Y_i) \sim N(0, \gamma)$

Scenario A2: $X_i \sim N(0, 1)$, $\epsilon_i | (X_i, Y_i) \sim N(0, \gamma_1 + \gamma_2 |X_i|)$

Scenario A3: $X_i \sim N(0, 1)$, $\epsilon_i | (X_i, Y_i) \sim N(0, \gamma_1 + \gamma_2 |X_i| + \gamma_3 |Y_i|)$

For each of the above scenarios, we examined two sets of measurement error parameters. In

Scenario A1, we took $\gamma = 0.5$ or 1 . In Scenarios A2 and A3, the two sets of measurement error parameters were chosen to make the unconditional variance of ϵ_i equal to about 0.5 or 1 , respectively. Scenario A1 is the classical additive error model, which is theoretically covered by RECS, SIMEX, and H&W (for N&S, the measurement error model assumption is satisfied but the entire function condition is not). Scenarios A2 and A3 involve heteroscedastic error models that are theoretically covered by RECS but not by N&S, SIMEX, or H&W.

Simulation Set B involved measurement error models of the form $W_i = X_i + \epsilon_i$, with non-normal ϵ_i . We considered two distributions for ϵ_i , the double-exponential distribution (DBLEXP(γ), with γ denoting the variance) and a modified chi-square distribution MODCHI which Huang and Wang (2001) used in their simulation work. Specifically, the MODCHI(γ) is defined to be the distribution of a χ_1^2 variate truncated at the value 5 , recentered to mean zero, and then rescaled to a variance of γ . In Scenario B3, we also take X to have a MODCHI distribution. The DBLEXP distribution is similar to the normal, but with heavier tails. The MODCHI distribution is highly skewed. The specific scenarios examined were as follows:

Scenario B1: $X_i \sim N(0, 1)$, $\epsilon_i | (X_i, Y_i) \sim \text{DBLEXP}(\gamma)$

Scenario B2: $X_i \sim N(0, 1)$, $\epsilon_i | (X_i, Y_i) \sim \text{MODCHI}(\gamma)$

Scenario B3: $X_i \sim \text{MODCHI}(1)$, $\epsilon_i | (X_i, Y_i) \sim \text{MODCHI}(\gamma)$

These scenarios are theoretically covered by RECS and H&W, but not by SIMEX or N&S.

For each of these scenarios, we ran simulations for $\gamma = 0.5$ and $\gamma = 1$. For the MODCHI distribution, integrals of the form $A_i g(x)$ were evaluated as described at the end of Web Appendix A.1. Note that the MODCHI(γ) is a non-regular distributional family: it has support that depends on γ . Hence, classical asymptotic theory for MLE's does not apply to the MLE of γ . However, due to the restricted range of γ values that are compatible with a given dataset due to the definition of the support, with an external validation sample of 70 the value of γ is estimated with virtually no error.

Simulation Set C considered the case of two error-prone covariates X_1, X_2 and one error-free covariate Z . For this simulation set, the sample size was 500, and the measurement error parameters were taken as known. For the RECS method, the basis function set was taken to be the tensor product of the univariate basis functions sets with $M = 6$, and in the quadrature calculations we took $K' = 20$ and $K = 10$. The true regression coefficients were $\beta_0 = \beta_1 = \beta_2 = \beta_3 = 1$. The scenarios examined were as follows.

Scenario C1: X_1, X_2, Z i.i.d. $N(0, 1)$; ϵ_1, ϵ_2 i.i.d. $N(0, \gamma)$, independent of X_1, X_2, Z

Scenario C2: X_1, X_2, Z i.i.d. $N(0, 1)$; ϵ_1, ϵ_2 conditionally independent given X_1, X_2, Z and distributed as $N(0, \gamma_1 + \gamma_2(|X_1| + |X_2| + |Z|))$

Scenario C3: $X_1, X_2 \sim \text{MODCHI}(1)$; $Z \sim N(0, 1)$; $\epsilon_1, \epsilon_2 \sim \text{MODCHI}(\gamma)$, all random variables independent of each other

In Scenarios C1 and C3, we took $\gamma = 1$, while in Scenario C2 we took $\gamma_1 = 0.4$ and $\gamma_2 = 0.25$, so that the unconditional variance of ϵ_1 and ϵ_2 was about 1. Scenario C1 is the classical additive error model, which is theoretically covered by RECS, SIMEX, and H&W (for N&S, the measurement error model assumption is satisfied but the entire function condition is not satisfied). Scenario C2 involves a heteroscedastic error model that is theoretically covered by RECS, but not by N&S, SIMEX, or H&W. Scenario C3 involves a non-normal error model that is theoretically covered by RECS and H&W, but not by SIMEX or N&S.

Finally, in Scenario D, we examined the effect of misspecifying the error distribution, in the setting of a single error-prone covariate. The details are described in Web Appendix C.2. The R code for the simulations can be found with the supplemental web materials or on the first author's website (<http://pluto.huji.ac.il/~mszucker>).

4.2 Simulation Results

Tables 1 and 2 present, respectively, the results of Simulation Studies A and B, while Table 3 presents the results of Simulation Study C. In Tables 1 and 2 we present the results only

for the slope parameter β_1 , which is the parameter of main interest. Results for the intercept parameter are given in the web appendix. In Table 3 we present results only for β_1 and β_3 , the slope parameters for X_1 and Z , respectively. The web appendix provides the results for β_0 and β_2 . The results for β_2 are similar to those for β_1 . The results of Simulation Scenario D are in the web appendix. Here we discuss the findings, focusing on estimation of the slope parameter. Overall, the RECS method performs very well in terms of bias and confidence interval coverage. Below we discuss how RECS compares with the competing methods.

[Table 1 about here.]

[Table 2 about here.]

[Table 3 about here.]

We begin with the Simulation Study A, the setting of normal measurement error with possible heteroscedasticity. RECS showed low bias throughout, especially when we look at the median of the estimates. N&S also generally showed low bias, although the bias was greater than that of RECS in a number of cases. SIMEX-NL also showed low bias in most cases (the last panel of Web Table 1 being a notable exception), but the estimation procedure was unsuccessful 6-10% of the time due to failure of the extrapolation process. SIMEX-L, SIMEX-Q, H&W, and MCMC performed markedly less well in terms of bias. The variability of the RECS estimates was comparable to that of the SIMEX-NL estimates and generally higher than that of the other methods.

We turn now to Simulation Study B, with non-normal measurement error. Again, RECS showed low bias throughout. N&S had low bias with double exponential error, and showed the best performance among the various methods in this case. With MODCHI error, RECS had low bias, while all the other methods had substantial bias.

We turn next to Simulation Study C, with two error-prone covariates and one error-free covariate. SIMEX and H&W performed poorly in estimating the slope parameters of the

two error-prone covariates. The SIMEX-NL method performed especially poorly, producing estimates way off in the wrong direction. The RECS method performed well. In Scenarios C1 (normal homoscedastic error) and C3 (MODCHI error), RECS performed markedly better than N&S in terms of producing estimates with low bias, while in Scenario C2 (normal heteroscedastic), the performance was similar, with RECS overestimating and N&S underestimating. With the MCMC method, the estimates with $\eta = 10$ were seriously biased, while the estimates with $\eta = 100$ had low to moderate bias.

In all three of the above scenarios, RECS generally yielded substantially better confidence interval coverage rates than the competing methods. Often the difference was substantial. We note that the SIMEX-NL method failed to admit a confidence interval in a considerable number of cases because the estimated variances were negative.

Finally, we turn to Simulation Study D, concerning the performance of RECS when the error model is misspecified (assumed normal but actually skewed normal or MODCHI). Aside from the case with both X and ϵ distributed MODCHI, the mean bias was in the range of 15-35% and the median bias was in the range 10-20%. Thus, in terms of median bias, RECS performed reasonably – certainly much better than the naive analysis with no measurement error correction. When both X and ϵ were distributed MODCHI, RECS performed poorly. However, this result is not too disturbing – because of the great difference between the MODCHI and normal distributions, it is unlikely that an analyst would mistakenly fit a normal model to MODCHI measurement errors. In principle, added robustness can be gained by using a flexible distributional form for the error model, such as the “semi-nonparametric normal” model of Gallant and Nychka (1987).

The degree of penalization in the RECS method tended to be very low throughout. The mean α value was less than 0.01 in all simulation scenarios studied.

5. Practical Illustration

We illustrate the method on data from the Nurses' Health Study (NHS). The NHS began in 1976 when 121,700 female nurses aged 30-55 returned a questionnaire about their lifestyle and their health. Here, we analyze the relationship between physical activity and mortality among women diagnosed with breast cancer during the NHS follow-up. This relationship was previously examined by Holmes et al. (2005). The present analysis involves a subset of the nurses included in the analysis of Holmes et al. Specifically, we consider the group of 1660 nurses who were diagnosed with breast cancer at least 10 years before the administrative end of the study, June 2002. This is the main study. The endpoint is the binary variable defined as breast cancer death within 10 years following diagnosis. The total number of such deaths was 188. In NHS, physical activity was assessed by a questionnaire in which women were asked how much time they spent on average during the past year on each of the most common forms of leisure time physical activity. The questionnaire results were then converted into metabolic equivalent task hours per week (METs). Validation data were available from 149 women from the NHS II study (Wolf et al., 1994), a study begun in 1989 which involved a cohort of U.S. female nurses similar to that of the NHS, and in which the same physical activity questions were asked. In our analysis, these data are regarded as arising from an external validation study. In the validation study, METs was assessed using both the questionnaire and a detailed activity diary, with the diary regarded as the gold standard. We denote the METs value based on diary data by X and the METs value based on questionnaire data by W . The degree of measurement error is considerable, with the correlation between X and W in the validation study being 0.47. Since the distribution of METs was skewed, we developed the measurement error model from the validation data using a transformed version of METs defined by $X^* = \log(1 + X)$. We considered two measurement error models, denoted MEM1 and MEM2. Both models were of the form $W_i^* = \omega_0 + \omega_1 X_i^* + \epsilon_i$. In MEM1,

ϵ_i was taken to have the $N(0, \sigma^2)$ distribution, independent of X_i^* . MEM2 incorporated dependence between ϵ_i and X_i^* , with the conditional distribution of ϵ_i given X_i^* taken to be $N(0, \gamma_1 + \gamma_2|X_i^* - \mu|)$, with $\mu = E[X_i^*]$. The parameters ω_0, ω_1 were estimated by simple linear regression in both models (weighted regression based on the MEM2 model for ϵ_i produced virtually identical estimates of ω_0 and ω_1). For MEM1, σ^2 was estimated by the regression MSE in the standard manner. For MEM2, γ_1 and γ_2 were estimated via regression analysis of the squares of the residuals obtained from MEM1; in this regression analysis, the t -test on γ_2 was borderline significant ($p=0.0614$), suggesting some evidence of heteroscedasticity. The estimates obtained were $\hat{\omega}_0 = 1.2271$, $\hat{\omega}_1 = 0.5653$, $\hat{\sigma}^2 = 0.8181$, $\hat{\gamma}_1 = 0.5883$, and $\hat{\gamma}_2 = 0.3497$. Graphical inspection of the normalized residuals based on the MEM2 model for ϵ_i showed reasonable conformity to a normal distribution. A preliminary analysis of the main study data indicated that the log odds of breast cancer could be reasonably expressed as a linear function of X^* , so we proceeded on this basis.

We ran RECS with 6 basis functions and $K = K' = 20$ quadrature points in the quadrature calculations. The relevant R code can be found with the supplemental web materials or on the first author's website. We also applied the SIMEX and N&S methods to the data. The H&W method is not relevant, since we have an external validation design rather than a replicate measures design. Standard errors for the RECS and N&S methods were computed using a sandwich-type estimator, while standard errors for SIMEX were computed using the jackknife method in the R package `simex`.

Table 4 presents the results for the various methods. The naive method was applied in two forms: (1) using W_i^* as is (Naive1) and (2) using $\tilde{W}_i^* = (W_i^* - \omega_0)/\omega_1$, thus correcting for location-scale bias but not for measurement error (Naive2). The table includes estimates of β_0 and β_1 and corresponding standard errors. The table also includes an estimate and a 95% confidence interval for the odds ratio associated with a change in X^* of 3.4, which

corresponds approximately to the difference between the extreme lower and upper METS categories defined in Holmes et al. (2005) (walking less than 1 hour per week vs. walking 8 or more hours per week). SIMEX-L yielded a slope estimate similar to that yielded by Naive2, while SIMEX-Q gave a slightly larger estimate. RECS-MEM1, N&S, and SIMEX-NL yielded slope estimates differing markedly from the Naive2 estimate, and the estimates yielded by these three methods were comparable. RECS-MEM2 yielded a slope estimate differing substantially from RECS-MEM1, showing the impact of a more refined error model.

[Table 4 about here.]

6. Discussion

We have presented a new “regularized corrected score” (RECS) approach to adjusting for covariate error in nonlinear regression problems. The approach builds on the corrected score method developed by Stefanski, Nakamura, and others. With a continuous error-prone covariate, the corrected score approach involves solving an integral equation. In many problems, an exact solution to this integral equation does not exist or cannot be practically obtained, and so we have proposed using an approximate solution obtained using the method of regularization. For logistic regression, a series of simulation studies showed that the method performs well in general, and offers some advantages over existing methods.

The RECS method yielded estimates with substantially less bias than competing estimators in the case of a single error-prone covariate with MODCHI error and in the case of two error-prone and one error-free covariates under two of the error models studied. The RECS estimator does tend to have higher variance than the other estimators, particularly the N&S estimator. We were unable to pinpoint the exact cause for the higher variance with the RECS method, but it is common for estimators that rely on weaker assumptions to be somewhat

more variable than estimators that rely on stronger assumptions. The RECS method had better confidence interval coverage rates than the competing methods in all scenarios studied.

We have developed the theory in the general setting of classical likelihood models, which covers, in particular, generalized linear models such as nonlinear regression, logistic regression, and Poisson regression. It is possible to extend the development to other settings. In particular, it is of interest to extend the method to Cox survival regression, using the work of Zucker and Spiegelman (2008) on corrected score analysis for the Cox model with a discrete error-prone covariate as a starting point. We plan to develop this extension in future work.

The computational complexity and load of the method is modest. For example, the data analysis described in the preceding section finished in about 1 minute of real time, when run in R in batch mode on a VMware virtual machine configured with one AMD 2700 MHz processor and 1GB memory, installed on a physical machine SUN FIRE X4240.

The method presented here is a functional method in the sense of not requiring information on the distribution of the true covariate. This is in contrast to many other measurement error methods, such as regression calibration and likelihood-based methods. We do rely on a parametric model for the conditional distribution of the surrogate variable given the true variable, but our simulations suggest that the performance of the estimates is robust to misspecification of parametric model unless the misspecification is extreme. Also, it is possible in principle to use a flexible parametric model such as Gallant and Nychka's (1987) "semi-nonparametric" model, which makes the reliance on parametric modeling less restrictive. The method is extremely general in terms of the types of measurement error models covered. It allows the measurement error to depend on the true covariate value and on other covariates. It also allows differential error, where the measurement error depends on the outcome. This flexibility is a distinct advantage relative to other methods.

ACKNOWLEDGEMENTS

This work has been supported by U.S. National Cancer Institute Grant 2R01CA050597. We thank Yijian (Eugene) Huang for providing code for the Huang and Wang (2001) method. We also thank the editor, associate editor, and referees for very helpful comments that led to substantial improvements in the paper.

SUPPLEMENTARY MATERIALS

Supplementary Materials are available with this paper at the Biometrics website on Wiley Online Library. These materials include Web Appendices A-C, referenced in Sections 3 and 4. Also included are an additional Web Appendix D with a brief overview of the computer codes used to generate the numerical results of this paper and a zip file with the codes themselves.

REFERENCES

- Akazawa, K., Kinukawa, N., and Nakamura, T. (1998). A note on the corrected score function corrected for misclassification. *Journal of the Japan Statistical Society* **28**:115–123.
- Box, G.E.P., and Tiao, G.C. (1973). *Bayesian Inference in Statistical Analysis*. Wiley: New York (reissued in paperback, 1992).
- Buzas, J. (2009). A note on corrected scores for logistic regression. *Statistics and Probability Letters* **79**:2351–2358
- Carroll, R.J., Ruppert, D., Stefanski, L.A., and Crainiceanu, C.M. (2006). *Measurement Error in Nonlinear Models: A Modern Perspective, 2nd ed.* Chapman and Hall / CRC: Boca Raton.
- Cook, J.R., and Stefanski, L.A. (1994). Simulation-extrapolation estimation in parametric measurement error models. *Journal of the American Statistical Association* **89**:1314–1328.

- Davidian, M., and Carroll, R.J. (1987) Variance function estimation. *Journal of the American Statistical Association* **82**:1079–1091.
- Delves, L.M., and Mohamed, J.L. (1985). *Computational Methods for Integral Equations*. Cambridge: Cambridge University Press.
- Devanarayan, V., and Stefanski, L.A. (2002). Empirical simulation extrapolation for measurement error models with replicate measurements. *Statistics and Probability Letters* **59**:219–225.
- Fuller, W.A. (1987). *Measurement Error Models*. Wiley: New York.
- Hansen, P.C. (1994). Regularization tools: a Matlab package for analysis and solution of discrete ill-posed problems. *Numerical Algorithms* **6**:1-35.
- Hansen, P.C. (2007). Regularization Tools Version 4.0 for Matlab 7.3. *Numerical Algorithms* **46**:189-194.
- Holmes, M.D., Chen, W.Y., Feskanich, D., Kroenke, C.H., and Colditz, G.A. (2005). Physical activity and survival after breast cancer diagnosis. *Journal of the American Medical Association* **293**:2479–2486.
- Huang, Y., and Wang, C.Y. (2001). Consistent function methods for logistic regression with errors in covariates. *Journal of the American Statistical Association* **95**:1209–1219.
- Gallant, A.R., and Nychka, D.W. (1987). Semiparametric maximum likelihood estimation. *Econometrica* **55**: 363-390.
- Kress, R. (1989). *Linear Integral Equations*. Berlin: Springer.
- Kong, F.H., and Gu, M. (1999). Consistent estimation in Cox proportional hazards model with covariate measurement errors. *Statistica Sinica* **9**:953-969.
- Lehmann, E.L. (1986). *Testing Statistical Hypotheses*, 2nd ed. Wadsworth and Brooks/Cole: Pacific Grove, CA.
- Nakamura, T. (1990). Corrected score function of errors-in-variables models: methodology

- and application to generalized linear models. *Biometrika* **77**:127–137.
- Nakamura, T. (1992). Proportional hazards model with covariates subject to measurement error. *Biometrics* **48**:829–838.
- Novick, S.J., and Stefanski, L.A. (2002). Corrected score estimation via complex variable simulation extrapolation. *Journal of the American Statistical Association* **97**:472–481.
- Richardson, S., Leblond, L., Jaussent, I., Green, P.J. (2002). Mixture models in measurement error problems, with reference to epidemiological studies. *Journal of the Royal Statistical Society, Series A* **165**: 549–566.
- Stefanski, L. (1989). Unbiased estimation of a nonlinear function of a normal mean with application to measurement-error models. *Communications in Statistics, Theory and Methods* **18**:4335–4358.
- Stefanski, L.A., Novick, S.J., and Devanarayan, V. (2005). Estimating a nonlinear function of a normal mean. *Biometrika* **92**:732–736.
- van der Vaart, A.W. (1998). *Asymptotic Statistics*. Cambridge: Cambridge University Press.
- Wolf, A.M., Hunter, D.J., Colditz, G.A., Manson, J.E., Stampfer, M.J., Corsano, K.A., Rosner, B., Krlska, A., and Willett, W.C. (1994) Reproducibility and validity of a self-administered physical activity questionnaire. *International Journal of Epidemiology* **23**:991–999.
- Zucker, D.M., and Spiegelman, D. (2008). Corrected score estimation in the proportional hazards model with misclassified discrete covariates. *Statistics in Medicine* **27**:1911–1933.

Received November 2011. Revised September 2012. Accepted October 2012.

Table 1
Simulation Study A
 $X \sim N(0, 1)$ and $\epsilon \sim N(0, \gamma_1 + \gamma_2|X| + \gamma_3Y)$
Results for Slope Parameter β_1 – Sample Size $n=200$, Validation Sample Size $m=70$

	M	MD	Emp-SD	IQ-SD	95% CI	F	M	MD	Emp-SD	IQ-SD	95% CI	F
	$\gamma_1 = 0.5, \gamma_2 = 0, \gamma_3 = 0$						$\gamma_1 = 1.0, \gamma_2 = 0, \gamma_3 = 0$					
Naive	0.6469	0.6379	0.1512	0.1520	0.360	0	0.4713	0.4660	0.1262	0.1264	0.027	0
RECS	1.0782	1.0314	0.3373	0.3013	0.958	0	1.1567	1.0231	0.5421	0.4266	0.933	2
N&S	1.0368	1.0257	0.2740	0.2867	0.907	0	0.9487	0.9859	0.2283	0.2365	0.897	8
SIMEX - Q	0.8940	0.8737	0.2245	0.2157	0.877	0	0.7182	0.6989	0.2048	0.2039	0.626	0
H&W	1.0600	0.9514	0.5264	0.3818	0.951	4	0.8840	0.7851	0.5249	0.3751	0.917	4
MCMC $\eta = 100$	1.2797	1.2364	0.4072	0.4324	0.950	0	1.2304	1.1671	0.4350	0.4848	0.970	0
	$\gamma_1 = 0.3, \gamma_2 = 0.25, \gamma_3 = 0$						$\gamma_1 = 0.7, \gamma_2 = 0.35, \gamma_3 = 0$					
Naive	0.6573	0.6494	0.1553	0.1573	0.390	0	0.4812	0.4749	0.1284	0.1279	0.036	0
RECS	1.0429	1.0054	0.3102	0.2784	0.949	0	1.0658	0.9797	0.4673	0.3663	0.912	1
N&S	1.0929	1.0881	0.3001	0.3208	0.899	1	0.9926	1.0305	0.2402	0.2348	0.890	7
SIMEX - Q	0.9169	0.8926	0.2368	0.2313	0.891	0	0.7358	0.7167	0.2115	0.2053	0.660	0
H&W	1.0370	0.9679	0.4451	0.2980	0.949	1	1.0310	0.9134	0.5109	0.3566	0.941	4
MCMC $\eta = 100$	1.2539	1.2292	0.3691	0.3750	0.950	0	1.2866	1.2396	0.4414	0.4678	0.940	0
	$\gamma_1 = 0.15, \gamma_2 = 0.25, \gamma_3 = 0.25$						$\gamma_1 = 0.35, \gamma_2 = 0.25, \gamma_3 = 0.50$					
Naive	0.6512	0.6437	0.1513	0.1488	0.373	0	0.5091	0.5029	0.1282	0.1285	0.060	0
RECS	1.0782	1.0316	0.3477	0.2921	0.950	0	1.1123	0.9989	0.4765	0.3793	0.929	0
N&S	1.1099	1.1037	0.2987	0.3215	0.896	2	1.0293	1.0674	0.2451	0.2514	0.858	12
SIMEX - Q	0.8718	0.8545	0.2241	0.2194	0.840	0	0.7259	0.7125	0.1986	0.1868	0.605	0
H&W	1.0410	0.9682	0.4373	0.3121	0.957	2	1.0600	0.9600	0.5004	0.3425	0.942	3
MCMC $\eta = 100$	1.0950	1.0748	0.3028	0.3490	0.990	0	1.1408	1.0934	0.3828	0.3606	0.980	0

N&S - Novick and Stefanski (2002); H&W - Huang and Wang (2001)

SIMEX results are based on $B = 100$, $\lambda = (0.5, 0.7, 0.9, 1.1, 1.3, 1.5, 1.7, 1.9)$ and the simex R library

M - empirical mean; MD - empirical median; Emp-SD - empirical standard deviation; IQ-SD - inter-quartile dispersion; 95% CI - empirical coverage rate of 95% Wald confidence interval (or HPD interval, for MCMC) ; F - number of samples with no solution

Table 2
Simulation Study B
Non-Normal Measurement Error with Normal or Non-Normal True Covariate
Results for Slope Parameter β_1 – Sample Size $n=200$, Validation Sample Size $m=70$

	M	MD	Emp-SD	IQ-SD	95% CI	F	M	MD	Emp-SD	IQ-SD	95% CI	F
$X \sim N(0, 1)$ and $\epsilon \sim \text{DBLEXP}(\gamma)$												
	$\gamma = 0.5$						$\gamma = 1.0$					
Naive	0.6378	0.6287	0.1609	0.1517	0.334	0	0.4659	0.4598	0.1370	0.1349	0.039	0
RECS	1.0684	0.9986	0.3611	0.3264	0.948	0	1.1362	1.0089	0.5714	0.4335	0.933	6
N&S	1.0471	1.0126	0.2970	0.3293	0.888	0	0.9625	1.0099	0.2794	0.2872	0.845	11
SIMEX - Q	0.8905	0.8641	0.2438	0.2276	0.871	0	0.7205	0.6970	0.2289	0.2165	0.629	0
H&W	1.0770	0.9848	0.4523	0.3202	0.951	0	0.9886	0.8985	0.4636	0.3336	0.942	1
MCMC $\eta = 100$	1.2987	1.2613	0.3796	0.4360	0.970	0	1.1476	1.0940	0.3785	0.4483	0.990	0
$X \sim N(0, 1)$ and $\epsilon \sim \text{MODCHI}(\gamma)$												
	$\gamma = 0.5$						$\gamma = 1.0$					
Naive	0.6822	0.6793	0.1706	0.1672	0.467	0	0.5196	0.5146	0.1547	0.1552	0.126	0
RECS	1.0555	1.0348	0.2981	0.2775	0.959	0	1.0869	1.0345	0.3970	0.3546	0.966	0
N&S	1.1449	1.1451	0.3230	0.3808	0.857	4	1.1232	1.1779	0.2711	0.2319	0.805	18
SIMEX - Q	0.9678	0.9500	0.2624	0.2617	0.890	0	0.8320	0.8234	0.2711	0.2765	0.747	0
H&W	1.0260	0.9809	0.3822	0.2861	0.956	2	0.9640	0.8949	0.4753	0.3069	0.927	3
MCMC $\eta = 100$	1.2154	1.1673	0.3372	0.2999	0.920	0	1.1502	1.1252	0.2996	0.2924	0.980	0
$X \sim \text{MODCHI}(1)$ and $\epsilon \sim \text{MODCHI}(\gamma)$												
	$\gamma = 0.5$						$\gamma = 1.0$					
Naive	0.5461	0.5348	0.1709	0.1662	0.250	0	0.3917	0.3824	0.1448	0.1402	0.063	0
RECS	1.1279	1.0479	0.4989	0.4415	0.938	0	1.1272	1.0319	0.5784	0.5316	0.932	0
N&S	1.2750	1.3552	0.4993	0.6506	0.672	51	1.0147	1.1508	0.4306	0.5522	0.545	212
SIMEX - Q	0.8175	0.7884	0.2870	0.2713	0.791	0	0.6396	0.6176	0.2602	0.2498	0.490	0
H&W	0.9610	0.8409	0.5507	0.3729	0.912	6	0.8084	0.6995	0.5894	0.4255	0.846	13
MCMC $\eta = 100$	1.0279	1.0105	0.4865	0.4735	0.970	0	0.8416	0.8245	0.4370	0.4278	0.930	0

N&S - Novick and Stefanski (2002); H&W - Huang and Wang (2001)

SIMEX results are based on $B = 100$, $\lambda = (0.5, 0.7, 0.9, 1.1, 1.3, 1.5, 1.7, 1.9)$ and the simex R library

M - empirical mean; MD - empirical median; Emp-SD - empirical standard deviation; IQ-SD - inter-quartile dispersion; 95% CI - empirical coverage rate of 95% Wald confidence interval (or HPD interval, for MCMC); F - number of samples with no solution

Table 3
Simulation Study C
Two Error-Prone Covariates and One Error-Free Covariate
Sample Size $n=500$

	β_1					β_3					F
	M	MD	Emp-SD	IQ-SD	95% CI	M	MD	Emp-SD	IQ-SD	95% CI	
Scenario C1: $X_1, X_2, Z \sim N(0, 1)$ and $\epsilon_1, \epsilon_2 \sim N(0, 1)$											
Naive	0.4296	0.4242	0.0804	0.0781	0.000	0.8626	0.8599	0.1201	0.1223	0.769	0
RECS	1.0817	1.0341	0.3116	0.2595	0.972	1.0525	1.0286	0.2079	0.1907	0.979	0
N&S	0.8286	0.8288	0.1316	0.1319	0.826	0.9483	0.9448	0.1382	0.1412	0.951	52
SIMEX - Q	0.8113	0.8038	0.1377	0.1333	0.571	0.9587	0.9524	0.1452	0.1519	0.920	0
H&W	0.6678	0.6675	0.5038	0.3360	0.800	1.0277	0.9679	0.4669	0.3039	0.918	14
MCMC $\eta=100$	1.0608	1.0335	0.2765	0.2508	0.870	1.0445	1.0418	0.1622	0.1834	0.970	0
Scenario C2: $X_1, X_2, Z \sim N(0, 1)$ and $\epsilon_1, \epsilon_2 \sim N(0, 0.4 + 0.25(X_1 + X_2 + Z))$											
Naive	0.4386	0.4334	0.0824	0.0792	0.001	0.8712	0.8694	0.1223	0.1223	0.791	0
RECS	1.1884	1.1122	0.3800	0.3126	0.986	1.0828	1.0482	0.2405	0.2007	0.983	1
N&S	0.8606	0.8619	0.1329	0.1312	0.842	0.9668	0.9655	0.1439	0.1465	0.966	46
SIMEX - Q	0.8345	0.8242	0.1431	0.1394	0.819	0.9793	0.9746	0.1516	0.1562	0.933	0
H&W	0.6165	0.6176	0.4809	0.3277	0.733	0.9627	0.9279	0.4248	0.2837	0.905	9
MCMC $\eta=100$	1.0338	1.0084	0.2709	0.2637	0.890	1.0358	1.0339	0.1626	0.1701	0.950	0
Scenario C3: $X_1, X_2 \sim \text{MODCHI}(1)$, $Z \sim N(0, 1)$, and $\epsilon_1, \epsilon_2 \sim \text{MODCHI}(1)$											
Naive	0.3853	0.3814	0.0887	0.0915	0.000	0.8883	0.8893	0.1206	0.1207	0.809	0
RECS	1.0430	0.9841	0.3726	0.3328	0.942	1.0413	1.0232	0.1856	0.1667	0.971	0
N&S	0.8656	0.8664	0.2576	0.2880	0.782	0.9261	0.9201	0.1396	0.1415	0.946	34
SIMEX - Q	0.7463	0.7315	0.1586	0.1598	0.445	0.9626	0.9588	0.1402	0.1399	0.915	0
H&W	0.4723	0.4276	0.2733	0.2108	0.654	0.9155	0.9023	0.1798	0.1638	0.971	1
MCMC $\eta=100$	1.1462	1.1199	0.3153	0.3259	0.910	1.0167	0.9887	0.1713	0.1519	0.910	0

N&S - Novick and Stefanski (2002); H&W - Huang and Wang (2001)

SIMEX results are based on $B = 100$, $\lambda = (0.5, 0.7, 0.9, 1.1, 1.3, 1.5, 1.7, 1.9)$ and the simex R library

M - empirical mean; MD - empirical median; Emp-SD - empirical standard deviation; IQ-SD - inter-quartile dispersion; 95% CI - empirical coverage rate of 95% Wald confidence interval (or HPD interval, for MCMC); F - number of samples with no solution

Table 4
NHS Results

Method	β_0		β_1		Odds Ratio for $\Delta X^* = 3.4$	
	Estimate	SE	Estimate	SE	Estimate	95% CI
Naive1	-1.7213	0.1436	-0.1802	0.0680	0.54	[0.34, 0.85]
Naive2	-1.9425	0.0862	-0.1019	0.0384	0.71	[0.55, 0.91]
RECS-MEM1	-1.7496	0.1328	-0.2733	0.1141	0.39	[0.18, 0.84]
RECS-MEM2	-1.6186	0.1693	-0.3970	0.1758	0.26	[0.08, 0.84]
N&S	-1.7501	0.1348	-0.2726	0.1161	0.40	[0.18, 0.86]
SIMEX-L	-1.9186	0.0887	-0.1223	0.0439	0.66	[0.49, 0.88]
SIMEX-Q	-1.8722	0.0936	-0.1630	0.0539	0.57	[0.40, 0.82]
SIMEX-NL	-1.7256	0.1155	-0.2966	0.0943	0.36	[0.19, 0.68]

Supplemental Web Materials

A Regularization Corrected Score Method for Nonlinear Regression Models with Covariate Error

David M. Zucker

Department of Statistics, Hebrew University

Mount Scopus, Jerusalem, Israel

email: mszucker@mssc.huji.ac.il

Malka Gorfine

Faculty of Industrial Engineering and Management

Technion - Israel Institute of Technology, Haifa, Israel

Yi Li

Department of Biostatistics

University of Michigan, Ann Arbor MI, USA

Mahlet G. Tadesse

Department of Mathematics and Statistics

Georgetown University, Washington DC, USA

and

Donna Spiegelman

Departments of Epidemiology and Biostatistics

Harvard School of Public Health, Boston MA, USA

Web Appendix A.1: Evaluation of Integrals of the Form $A_i g(x)$

As noted in the main text, the RECS procedure requires evaluation of integrals of the form

$$E[g(W_i)|X_i = x] = A_i g(x) = \int a_i(x, w)g(w)dw. \quad (1)$$

In our simulation studies, we work with measurement error models of the form $W = X + \sigma(X)\varepsilon$, where ε is a random variable with density f_ε , independent of X . In this case, we can write

$$A_i g(x) = \int f_\varepsilon(v)g(x + \sigma(x)v)dv.$$

This integral can be evaluated using a quadrature approximation of the form

$$A_i g(x) \doteq \sum_{k=1}^{K'} r_k g(x + \sigma(x)v_k)$$

for suitable quadrature points and weights v_k and r_k . One case we consider is $\varepsilon \sim N(0, 1)$, and in this case we use (modified) Gauss-Hermite points and weights. Another case we consider is the case where ε is the double-exponential (Laplace) distribution with variance 1, and in this case we use Gauss-Laguerre type points and weights (with the points and weights on the negative side of the real axis being the mirror image of those on the positive side).

In cases where specialized methods of the above sort are not readily available, a more general numerical approach can be used. In particular, if we let $F_{W|X}$ be the conditional distribution function of W given X , then $A_i g(x)$ can be evaluated as

$$A_i g(x) \doteq \frac{1}{K'} \sum_{k=1}^{K'} g(F_{W|X}^{-1}(v_k|x))$$

with $v_k = (k - \frac{1}{2})/K'$.

For the MODCHI distribution used in the simulation study, integrals of the form

$A_i g(x)$ were evaluated using the representation

$$A_i g(x) = \frac{2}{\Pr(\chi_1^2 \leq 5)} \int_0^{\sqrt{5}} \varphi(v) g \left(x + \gamma \left[\frac{v^2 - \mu_{MC}}{\sigma_{MC}} \right] \right) dv$$

in conjunction with Gauss-Hermite quadrature, where μ_{MC} and σ_{MC} denote, respectively, the mean of the chi-square distribution truncated at 5.

Web Appendix A.2: Theorem of Section 3.2 - Regularity Conditions and Proof

Regularity Conditions

A1. The parameter space Θ is compact with a nonempty interior which includes the true value θ_0 .

A2. The function $\mathbf{u}_i(x, \theta)$ is continuously differentiable in θ over Θ for every x , with derivative that is bounded over x by an L^2 function of x .

A3. The matrix $\mathbf{D}_E(\theta)$ defined by $(D_E)_{rs}(\theta) = -E[(\partial^2 / \partial \theta_r \partial \theta_s) \log f(Y|X, \mathbf{Z}, \theta)]$ is positive definite over Θ .

A4. The null space $\mathcal{N}(A_i^*)$ consists only of the zero function, i.e., the only solution to $A_i^* h = 0$ is $h = 0$.

Proof

In proving Part (a) of the theorem, we rely on the L^2 theory of integral operators as set forth, for example, in Kress (1989). We recall that, for a general integral operator B , the range of B is defined as $\mathcal{R}(B) = \{h \in L^2 : h = Bg \text{ for some } g \in L^2\}$ and the null space of B is defined $\mathcal{N}(B) = \{g \in L^2 : Bg = 0\}$. We use a superscript \perp to denote orthogonal complement and the notation $cl(C)$ to denote the L^2 closure of a set $C \subset L^2$.

Kress (1989, Theorem 15.8) states that for a bounded linear operator $B : L^2 \rightarrow L^2$ with adjoint B^* , we have $\mathcal{N}(B^*)^\perp = cl(\mathcal{R}(B))$. Since $a_i(x, w)$ is a conditional density,

the operator A_i is a bounded linear operator with norm 1. Assumption A4 specifies that $\mathcal{N}(A_i^*)$ consists only of the zero function. It follows that the L^2 closure of $\mathcal{R}(A_i)$ is equal to the whole of L^2 . This, in turn, implies that $\inf_{\bar{\delta} \in L^2} \|A_i \bar{\delta} - g\| = 0 \forall g \in L^2$, although the infimum is not necessarily attained, which, in our setting, corresponds to the fact that an exact corrected score may not exist.

Now, for a given L^2 function δ , let $\bar{\delta}(\alpha)$ denote the minimizer of $\mathcal{L}(\bar{\delta}; A_i, \delta, \alpha) = \|A_i \bar{\delta} - \delta\|^2 + \alpha \|\bar{\delta}\|^2$. We claim that $\lim_{\alpha \rightarrow 0} \|\bar{\delta}(\alpha) - \delta\| = 0$. The proof is simple, and is implicit in Kress (1989, Chapter 16), but we give it for completeness. We have

$$\|A_i \bar{\delta}(\alpha) - \delta\|^2 \leq \|A_i \bar{\delta}(\alpha) - \delta\|^2 + \alpha \|\bar{\delta}(\alpha)\|^2 \leq \|A_i g - \delta\|^2 + \alpha \|g\|^2$$

for any $g \in L^2$. Letting $\alpha \rightarrow 0$, we get $\lim_{\alpha \rightarrow 0} \|A_i \bar{\delta}(\alpha) - \delta\|^2 \leq \|A_i g - \delta\|^2$. Since g was arbitrary, we get $\lim_{\alpha \rightarrow 0} \|A_i \bar{\delta}(\alpha) - \delta\|^2 \leq \inf_g \|A_i g - \delta\|^2$, but the infimum on the right side, as we just saw, is equal to zero. We have thus proved the claim.

In the context of our RECS estimator, we obtain the following result: defining $r_{ij}(x, \boldsymbol{\theta}, \alpha) = A_i \bar{u}_{ij}(x, \boldsymbol{\theta}, \alpha) - u_{ij}(x, \boldsymbol{\theta})$, we have $\|r_{ij}(\cdot, \boldsymbol{\theta}, \alpha)\| \rightarrow 0$ as $\alpha \rightarrow 0$. At this point, we have this convergence only at a fixed value of $\boldsymbol{\theta}$. However, since Θ is assumed compact and $u_{ij}(x, \boldsymbol{\theta})$ is continuous in $\boldsymbol{\theta}$, pointwise convergence implies uniform convergence. This yields the desired result.

Let us now turn to Part (b). Define $\Delta_{ijs}(x) = u_{ijs}(x, \boldsymbol{\theta}) - A_i u_{ijs}(x, \boldsymbol{\theta})$ and $\bar{\Delta}_{ijs}(w, \boldsymbol{\theta}, \alpha) = \bar{u}_{ijs}(w, \boldsymbol{\theta}, \alpha) - u_{ijs}(w, \boldsymbol{\theta})$. As stated near the end of Section 2 of the main paper, the minimizer of $\mathcal{L}(\bar{\delta}; A_i, \delta, \alpha)$ can be expressed as $\bar{\delta} = (A_i^* A_i + \alpha \mathcal{I})^{-1} A_i^* \delta$. This is a linear function of δ . It follows that, just as the function $\bar{\Delta}_{ij}(\cdot, \boldsymbol{\theta}, \alpha)$ is the minimizer of $\|A_i \bar{\delta} - \Delta_{ij}\|^2 + \alpha \|\bar{\delta}\|^2$, so, too, the function $\bar{\Delta}_{ijs}(\cdot, \boldsymbol{\theta}, \alpha)$ is the minimizer of $\|A_i \bar{\delta} - \Delta_{ijs}\|^2 + \alpha \|\bar{\delta}\|^2$. We can therefore apply the arguments just used in the proof of Part (a) to prove Part (b).

Given the results in Parts (a) and (b) of the theorem, the results in Parts (c) and

(e) follow from standard estimating equations theory, as in, for example, Huber (1967), White (1982), and van der Vaart (1998, Ch. 5).

Finally, we turn to Part (d). From the development in Part (a), we have

$$\bar{\mathbf{u}}_E(\tilde{\boldsymbol{\theta}}^{(\alpha)}, \alpha) = E[\bar{\mathbf{u}}_i(W_i, \tilde{\boldsymbol{\theta}}^{(\alpha)})] = E[\mathbf{u}_i(X_i, \tilde{\boldsymbol{\theta}}^{(\alpha)})] + E[\mathbf{r}_i(X_i, \tilde{\boldsymbol{\theta}}^{(\alpha)}, \alpha)].$$

The left side of the above equation, by definition, is zero. Expanding the first term of the right side in a first-order Taylor series around $\boldsymbol{\theta}_0$, we get

$$-\mathbf{D}_E(\boldsymbol{\theta}^\#)(\tilde{\boldsymbol{\theta}}^{(\alpha)} - \boldsymbol{\theta}_0) + E[\mathbf{r}_i(X_i, \tilde{\boldsymbol{\theta}}^{(\alpha)}, \alpha)] = 0.$$

where $\mathbf{D}_E(\boldsymbol{\theta})$ is as defined in Assumption A3 and $\boldsymbol{\theta}^\#$ is some value between $\boldsymbol{\theta}_0$ and $\tilde{\boldsymbol{\theta}}^{(\alpha)}$. Part (a) implies that the second term on the right side tends to zero as $\alpha \rightarrow 0$. Given this, along with the nonsingularity condition A3, we obtain the desired conclusion $\tilde{\boldsymbol{\theta}}^{(\alpha)} \rightarrow \boldsymbol{\theta}_0$ as $\alpha \rightarrow 0$.

Web Appendix A.3: Correcting the Covariance for Estimation of $\mathbf{A}^{(i)}$

We describe here how to correct the covariance of $\hat{\boldsymbol{\theta}}$ for estimation of $\mathbf{A}^{(i)}$. In the development below, we generally suppress the dependence of various quantities on the penalty parameter α .

The parameter $\boldsymbol{\xi}$ entering into $a_i(x, w, \boldsymbol{\xi})$ is estimated on the basis of an external or internal validation sample, or a replicate measures study, of size denoted by m . Let $\boldsymbol{\xi}_0$ denote the true value of $\boldsymbol{\xi}$. We assume that the estimator $\hat{\boldsymbol{\xi}}$ has an approximate normal distribution with mean $\boldsymbol{\xi}_0$ and covariance matrix $m^{-1}\boldsymbol{\Gamma}$, along with an estimator of the matrix $\boldsymbol{\Gamma}$. This setup is a typical one in practice. For the asymptotics we assume that m and n are of the same order of magnitude, i.e., $m/n \rightarrow \zeta$ for some constant ζ as $n \rightarrow \infty$. Otherwise the error in $\hat{\boldsymbol{\xi}}$ will either be dominated by or will dominate the error in $\boldsymbol{\theta}$ due to the variation in the outcome data. Typically ζ will be between 0 and 1. The asymptotic

covariance matrix of $\boldsymbol{\xi}$ may then be expressed as $n^{-1}\zeta^{-1}\boldsymbol{\Gamma}$. To emphasize the dependence of the corrected score on $\boldsymbol{\xi}$, we write $\bar{\mathbf{U}}(\boldsymbol{\theta}, \boldsymbol{\xi})$. The estimated asymptotic covariance matrix of $\bar{\mathbf{U}}(\tilde{\boldsymbol{\theta}}^{(\alpha)}, \boldsymbol{\xi}_0)$ is $n^{-1}\mathbf{F}^{(\alpha)}(\hat{\boldsymbol{\theta}})$ where $\mathbf{F}^{(\alpha)}(\boldsymbol{\theta})$ is as defined in (8). We denote the asymptotic covariance between $\bar{\mathbf{U}}(\boldsymbol{\theta})$ and $\hat{\boldsymbol{\xi}}$ by $n^{-1}\boldsymbol{\Upsilon}$. The form of $\boldsymbol{\Upsilon}$ depends on the type of data used to estimate $\boldsymbol{\xi}$, and will be discussed shortly.

Let $\bar{\mathbf{U}}'(\boldsymbol{\theta}, \boldsymbol{\xi})$ denote the matrix whose (r, ν) element is the partial derivative of $\bar{U}_r(\boldsymbol{\theta}, \boldsymbol{\xi})$ with respect to ξ_ν . By Taylor expansion, we have

$$\mathbf{0} = \bar{\mathbf{U}}(\hat{\boldsymbol{\theta}}^{(\alpha)}, \hat{\boldsymbol{\xi}}) = \bar{\mathbf{U}}(\tilde{\boldsymbol{\theta}}^{(\alpha)}, \boldsymbol{\xi}_0)\bar{\mathbf{D}}(\tilde{\boldsymbol{\theta}}^{(\alpha)}, \boldsymbol{\xi}_0)(\hat{\boldsymbol{\theta}}^{(\alpha)} - \tilde{\boldsymbol{\theta}}^{(\alpha)}) + \bar{\mathbf{U}}'(\tilde{\boldsymbol{\theta}}^{(\alpha)}, \boldsymbol{\xi}_0)(\hat{\boldsymbol{\xi}} - \boldsymbol{\xi}_0) + o_p(1/n),$$

leading to

$$-\sqrt{n}(\hat{\boldsymbol{\theta}}^{(\alpha)} - \tilde{\boldsymbol{\theta}}^{(\alpha)}) = \bar{\mathbf{D}}(\tilde{\boldsymbol{\theta}}^{(\alpha)}, \boldsymbol{\xi}_0)^{-1} \left[\{\sqrt{n}\bar{\mathbf{U}}(\tilde{\boldsymbol{\theta}}^{(\alpha)}, \boldsymbol{\xi}_0)\} + \bar{\mathbf{U}}'(\tilde{\boldsymbol{\theta}}^{(\alpha)}, \boldsymbol{\xi}_0)\{\sqrt{n}(\hat{\boldsymbol{\xi}} - \boldsymbol{\xi}_0)\} \right] + o_p(1/n).$$

Accordingly, the estimated asymptotic covariance matrix of $\sqrt{n}(\hat{\boldsymbol{\theta}} - \tilde{\boldsymbol{\theta}}^{(\alpha)})$ is $\hat{\mathbf{V}}(\hat{\boldsymbol{\theta}}, \hat{\boldsymbol{\xi}})$, where $\hat{\mathbf{V}}(\boldsymbol{\theta}, \boldsymbol{\xi})$ is now defined as

$$\hat{\mathbf{V}}(\boldsymbol{\theta}, \boldsymbol{\xi}) = \bar{\mathbf{D}}(\boldsymbol{\theta}, \boldsymbol{\xi})^{-1} \left[\mathbf{F}^{(\alpha)}(\boldsymbol{\theta}) + \zeta^{-1}\bar{\mathbf{U}}'(\boldsymbol{\theta}, \boldsymbol{\xi})\hat{\boldsymbol{\Gamma}}\bar{\mathbf{U}}'(\boldsymbol{\theta}, \boldsymbol{\xi})^T + \hat{\boldsymbol{\Upsilon}}\bar{\mathbf{U}}'(\boldsymbol{\theta}, \boldsymbol{\xi})^T \right] \bar{\mathbf{D}}(\boldsymbol{\theta}, \boldsymbol{\xi})^{-1}.$$

If $\boldsymbol{\xi}$ is estimated from an external validation or replicate measures study, then $\hat{\boldsymbol{\xi}}$ is obviously independent of the corrected score function $\bar{\mathbf{U}}(\boldsymbol{\theta})$, and thus $\boldsymbol{\Upsilon} = \mathbf{0}$. When $\boldsymbol{\xi}$ is estimated from an internal validation or replicate measures study, $\boldsymbol{\Upsilon}$ is nonzero and must be estimated. We consider the setting where the validation/replicate data are i.i.d. across individuals, and $\boldsymbol{\xi}$ is estimated by maximum likelihood. Let $g_i(\boldsymbol{\xi})$ denote the log likelihood function for the validation/replicate data on individual i . The overall normalized log likelihood for $\boldsymbol{\xi}$ is then $g(\boldsymbol{\xi}) = m^{-1}\sum_{i \in \mathcal{R}} g_i(\boldsymbol{\xi})$, where \mathcal{R} denotes the set of individuals in the internal validation/replicate sample. Let $\mathbf{g}'(\boldsymbol{\xi})$ and $\mathbf{g}''(\boldsymbol{\xi})$ denote the gradient vector and Hessian matrix, respectively, of $g(\boldsymbol{\xi})$, and let $\mathbf{g}'_i(\boldsymbol{\xi})$ denote the gradient of $g_i(\boldsymbol{\xi})$. We can then express $\hat{\boldsymbol{\xi}}$ in terms of the classic asymptotic approximation $\hat{\boldsymbol{\xi}} \doteq -\mathbf{g}''(\boldsymbol{\xi}_0)^{-1}\mathbf{g}'(\boldsymbol{\xi}_0)$.

Define $\mathbf{\Omega} = \text{Cov}(\bar{\mathbf{u}}_i(\boldsymbol{\theta}_0, \boldsymbol{\xi}_0), \mathbf{g}'_i(\boldsymbol{\xi}_0))$. The matrix $\mathbf{\Omega}$ can be estimated empirically by

$$\hat{\mathbf{\Omega}} = \frac{1}{m} \sum_{i \in \mathcal{R}} \bar{\mathbf{u}}_i(\hat{\boldsymbol{\theta}}, \hat{\boldsymbol{\xi}}) \mathbf{g}'_i(\hat{\boldsymbol{\xi}})^T. \quad (2)$$

We then estimate $\mathbf{\Upsilon}$ by $\hat{\mathbf{\Upsilon}} = \hat{\mathbf{\Omega}} \mathbf{g}''(\hat{\boldsymbol{\xi}})^{-1}$. In addition, in the present setup we have $\hat{\mathbf{\Gamma}} = \mathbf{g}''(\hat{\boldsymbol{\xi}})^{-1}$.

In principle, expressions can be worked out for the partial derivatives that make up the matrix $\bar{\mathbf{U}}'(\boldsymbol{\theta}, \boldsymbol{\xi})$, but the algebra is cumbersome. Therefore, in our practical implementation, we use numerical partial derivatives.

Web Appendix B: Description of Bayesian MCMC Method

Richardson et al. (2002) proposed a semiparametric Bayesian measurement error method that uses a mixture of normal distributions to model the distribution of the error prone covariates X . They considered the case in which the main study sample consists of n units on which only W and Y are observed and a validation sample of size m is available with observations on X, W, Y . All $n^* = n + m$ observations are used to formulate a hierarchical model for the measurement error model, as follows ($i = 1, \dots, n^*$):

$$\begin{aligned} Y_i | X_i, \beta_0, \beta_1 &\sim \text{Bernoulli}(p_i) \quad \text{with } p_i = \frac{\exp(\beta_0 + \beta_1 X_i)}{1 + \exp(\beta_0 + \beta_1 X_i)} \\ W_i | X_i, \alpha_0, \alpha_1, \gamma &\sim N(\alpha_0 + \alpha_1 X_i, \gamma) \\ X_i | \boldsymbol{\omega}, \boldsymbol{\mu}, \boldsymbol{\sigma}^2, k &\sim \sum_{j=1}^k \omega_j N(\mu_j, \sigma_j^2). \end{aligned}$$

Conjugate priors are specified for the model parameters with hyperparameters defined to lead to weakly informative priors centered around estimates from the validation group. Both the validation set and the main study contribute to the joint posterior distribution

which is given by:

$$p(\boldsymbol{\beta}) p(\boldsymbol{\alpha}) p(\gamma) p(\boldsymbol{\omega}|k) p(\boldsymbol{\mu}|k) p(\boldsymbol{\sigma}^2|k) p(k) \times \prod_{i=1}^{n^*} [p(Y_i|X_i, \boldsymbol{\beta}) p(W_i|X_i, \boldsymbol{\alpha}, \gamma) p(X_i|c_i, \boldsymbol{\mu}, \boldsymbol{\sigma}^2) p(c_i|\boldsymbol{\omega}, k)], \quad (3)$$

where c_i is the component allocation indicator for X_i . The model is fit using a Markov chain Monte Carlo (MCMC) algorithm that iterates over the following steps:

1. update the regression coefficients and measurement error parameters using a Metropolis-Hastings algorithm
2. update each X_i ($i = 1, \dots, n$) in the main study using a random walk Metropolis step
3. update each of the mixture component parameters using Gibbs sampling
4. update the number of mixture components, k , and consequently the relevant mixture parameters using split/merge and birth/death moves.

In the scenarios we are considering, the validation sample includes data on (X, W) only and not on Y (for Scenario A3 [differential error], the measurement error parameters were fixed at their true values and assumed known in the MCMC analysis). In addition, we consider measurement error models of the form $W_i = X_i + \epsilon_i$ with different parametric distributions for ϵ_i as described in the various simulation scenarios. Some slight modifications in the Richardson et al. (2002) method are required to accommodate these features. In particular, an additional step in the MCMC algorithm is needed to update the unobserved Y_i ($i = 1, \dots, m$) in the validation sample. In addition, the prior settings for the regression coefficients need to be modified. Since we can no longer obtain regression coefficient estimates from the validation set, we center the prior for β_j around the naive estimates $\tilde{\beta}_j$ ($j = 0, 1$). An important implication of not observing Y in the validation

sample is that the prior for the β_j needs to be relatively informative; we set the prior variance to be proportional to the variance of the naive estimate. For the updates of the measurement error parameters and the unobserved X_i 's, the assumed parametric form of the measurement error model is used in the corresponding MCMC steps. The prior specifications we used are as follows:

$$\begin{aligned} \beta &\sim N(\tilde{\beta}, \eta \widehat{\text{Var}}(\tilde{\beta})) \\ k &\sim \text{Truncated-Poisson}(\lambda = 3) \quad k = 1, \dots, 30 \\ c_i|k &\sim \text{Multinomial}(1; \omega_1, \dots, \omega_k) \quad \omega \sim \text{Dirichlet}(1, \dots, 1) \\ \mu_j &\sim N(\xi, R^2) \quad \sigma_j^{-2} \sim \text{Gamma}(2, b_\sigma) \quad b_\sigma \sim \text{Gamma}(0.2, 10/R^2), \end{aligned}$$

where ξ and R are respectively set to the midpoint and the range of the initial values for X_i , taken to be $\hat{X}_i \sim N(W_i, \widehat{\text{Var}}(\hat{\epsilon}_i))$ with $\widehat{\text{Var}}(\hat{\epsilon}_i)$ estimated according to the assumed error model. The hyperparameter η controls the strength of the prior information for the regression coefficients β_j with larger values leading to less informative priors. In our analyses, we consider $\eta = 10$ and 100 .

Web Appendix C.1: Additional Results for Simulation Scenarios A-C

Tables 1, 2, and 3-5 provide results for Simulation Scenarios A, B, and C, respectively. These tables include the results provided in the main paper along with additional results beyond those provided in the main paper. For Scenario C we add results on the slope β_2 of the second error-prone covariate. For all scenarios we add results on the intercept β_0 and results for the SIMEX-L method, the SIMEX-NL method, and the MCMC method with $\eta = 10$.

Web Appendix C.2: Simulation Scenario D

Simulation Set D examined, in the setting of Simulation Sets A and B, the effect of misspecifying the error distribution. We generated the errors according to one of two

possible non-normal distributions, but implemented our method assuming the errors are normal. The non-normal error distributions used were the MODCHI(γ) distribution and a modified version of Azzalini's (1985) skewed normal distribution. Azzalini's skewed normal distribution $SN(\lambda)$ has density $2\phi(y)\Phi(\lambda y)$, $y \in \mathbb{R}$, where ϕ and Φ denote the standard normal density and distribution function, respectively, and λ is a parameter that regulates the skewness ($\lambda = 0$ gives the standard normal). Our modified version recenters to mean zero and then rescales to the specified variance. We took $W_i = X_i + \epsilon_i$, with X_i taken to be either $N(0, 1)$, $SN(50)$, or MODCHI, and the distribution of ϵ_i taken to be either the MODCHI(γ) or the the modified $SN(\lambda)$ with $\lambda = 50$. The skewness of the $SN(50)$ distribution is 1, and that of the MODCHI is 1.7.

Web Appendix D: Computer Code

The computer code for the numerical results presented in this paper are contained in a zip file called RECS Paper Codes.zip. This file contains five folders labeled as follows: RECS Method, N-S Method, SIMEX Method, H-W Method, and MCMC Method. In addition, it contains a file called `readme.txt` with some explanatory notes.

The RECS Method folder contains three zip files with the codes for the RECS method, as follows:

R code for Simulation Studies A, B, and D: `recs-siml-univ.zip`

R code for Simulation Study C: `recs-siml-multiv.zip`

R code for running the example: `recs-example.zip`

The codes were written in R, with some subroutines written in Fortran to increase the speed of the computation. The codes are also available on the first author's website

<http://pluto.huji.ac.il/~mszucker>

under the same file names (add `/recs-siml-univ.zip` to the URL for the first file, and similarly for the other two). In the event that the codes are updated, the updated versions

will be posted on the above website. The other folders contain the codes for the other methods. More details are provided in the `readme.txt` file.

ADDITIONAL REFERENCES

Huber, P. (1967). The behavior of maximum likelihood estimates under nonstandard conditions. In *Proceedings of the Fifth Berkeley Symposium on Mathematical Statistics and Probability*, Vol. 1, pp. 221–233. University of California Press: Berkeley.

van der Vaart, A.W. (1998). *Asymptotic Statistics*. Cambridge University Press: Cambridge.

White, H. (1982). Maximum likelihood estimation of misspecified models. *Econometrica* **50**:1–25.

Web Table 1

Simulation Study A

$$X \sim N(0, 1) \text{ and } \epsilon \sim N(0, \gamma_1 + \gamma_2|X| + \gamma_3Y)$$

Sample Size $n=200$, Validation Sample Size $m=70$

	β_0					β_1					F	
	M	MD	Emp-SD	IQ-SD	95% CI	M	MD	Emp-SD	IQ-SD	95% CI		
				$\gamma_1 = 0.5$	$\gamma_2 = 0$	$\gamma_3 = 0$						
Naive	0.9526	0.9459	0.1737	0.1695	0.930	0.6469	0.6379	0.1512	0.1520	0.360	0	
RECS	1.0323	1.0217	0.2122	0.1983	0.959	1.0782	1.0314	0.3373	0.3013	0.958	0	
N&S	1.0085	1.0004	0.1902	0.1881	0.958	1.0368	1.0257	0.2740	0.2867	0.907	0	
SIMEX - Q	0.9942	0.9840	0.1890	0.1890	0.953	0.8940	0.8737	0.2245	0.2157	0.877	0	
SIMEX - L	0.9718	0.9620	0.1800	0.1764	0.942	0.7619	0.7501	0.1803	0.1794	0.663	0	
SIMEX - NL	1.0548	1.0252	0.3080	0.2261	0.934	1.0681	1.0161	0.3312	0.3032	0.977	61	
H&W	0.9470	0.9752	0.6279	0.2387	0.975	1.0600	0.9514	0.5264	0.3818	0.951	4	
MCMC $\eta = 10$	1.0209	1.0159	0.1965	0.2011	0.93	1.0278	1.0194	0.2635	0.2867	0.970	0	
MCMC $\eta = 100$	1.1000	1.1028	0.2299	0.2360	0.96	1.2797	1.2364	0.4072	0.4324	0.950	0	
				$\gamma_1 = 1$	$\gamma_2 = 0$	$\gamma_3 = 0$						
Naive	0.9231	0.9153	0.1694	0.1665	0.909	0.4713	0.4660	0.1262	0.1264	0.027	0	
RECS	1.0613	1.0346	0.2683	0.2283	0.963	1.1567	1.0231	0.5421	0.4266	0.933	2	
N&S	0.9835	0.9774	0.1873	0.1926	0.942	0.9487	0.9859	0.2283	0.2365	0.897	8	
SIMEX - Q	0.9645	0.9515	0.1855	0.1809	0.932	0.7182	0.6989	0.2048	0.2039	0.626	0	
SIMEX - L	0.9390	0.9270	0.1748	0.1675	0.922	0.5663	0.5595	0.1525	0.1534	0.150	0	
SIMEX - NL	1.0564	1.0068	0.4722	0.2328	0.799	1.1431	0.9997	0.5593	0.4151	0.939	63	
H&W	0.9065	0.9495	0.6558	0.2483	0.975	0.8840	0.7851	0.5249	0.3751	0.917	4	
MCMC $\eta = 10$	0.9802	0.9705	0.1899	0.1939	0.95	0.8551	0.8296	0.2428	0.2878	0.870	0	
MCMC $\eta = 100$	1.0910	1.0840	0.2366	0.2477	0.95	1.2304	1.1671	0.4350	0.4848	0.970	0	
				$\gamma_1 = 0.3$	$\gamma_2 = 0.25$	$\gamma_3 = 0$						
Naive	0.9522	0.9446	0.1734	0.1728	0.932	0.6573	0.6494	0.1553	0.1573	0.390	0	
RECS	1.0233	1.0121	0.2057	0.1953	0.958	1.0429	1.0054	0.3102	0.2784	0.949	0	
N&S	1.0050	0.9963	0.1861	0.1827	0.958	1.0929	1.0881	0.3001	0.3208	0.899	1	
SIMEX - Q	0.9933	0.9825	0.1884	0.1913	0.950	0.9169	0.8926	0.2368	0.2313	0.891	0	
SIMEX - L	0.9713	0.9615	0.1796	0.1779	0.943	0.7750	0.7620	0.1857	0.1868	0.682	0	
SIMEX - NL	1.0427	1.0199	0.2912	0.2209	0.932	1.1265	1.0607	0.3867	0.3321	0.972	74	
H&W	0.9408	0.9680	0.5539	0.2150	0.973	1.0370	0.9679	0.4451	0.2980	0.949	1	
MCMC $\eta = 10$	1.0415	1.0487	0.1988	0.1990	0.93	1.0953	1.0994	0.2649	0.2854	0.970	0	
MCMC $\eta = 100$	1.0946	1.1061	0.2189	0.2245	0.95	1.2539	1.2292	0.3691	0.3750	0.950	0	
				$\gamma_1 = 0.7$	$\gamma_2 = 0.35$	$\gamma_3 = 0$						
Naive	0.9238	0.9145	0.1695	0.1664	0.905	0.4812	0.4749	0.1284	0.1279	0.036	0	
RECS	1.0363	1.0161	0.2423	0.2153	0.960	1.0658	0.9797	0.4673	0.3663	0.912	1	
N&S	0.9763	0.9696	0.1845	0.1875	0.941	0.9926	1.0305	0.2402	0.2348	0.890	7	
SIMEX - Q	0.9650	0.9536	0.1855	0.1861	0.933	0.7358	0.7167	0.2115	0.2053	0.660	0	
SIMEX - L	0.9397	0.9285	0.1748	0.1668	0.919	0.5780	0.5685	0.1552	0.1542	0.177	0	
SIMEX - NL	1.0484	1.0065	0.4375	0.2305	0.807	1.2113	1.0383	0.6183	0.4470	0.955	63	
H&W	0.9388	0.9663	0.6524	0.2431	0.972	1.0310	0.9134	0.5109	0.3566	0.941	4	
MCMC $\eta = 10$	1.0035	0.9925	0.1937	0.2040	0.94	0.9550	0.9292	0.2552	0.2945	0.940	0	
MCMC $\eta = 100$	1.1025	1.1115	0.2345	0.2208	0.96	1.2866	1.2396	0.4414	0.4678	0.940	0	
				$\gamma_1 = 0.15$	$\gamma_2 = 0.25$	$\gamma_3 = 0.25$						
Naive	0.9524	0.9433	0.1715	0.1717	0.935	0.6512	0.6437	0.1513	0.1488	0.373	0	
RECS	1.0274	1.0114	0.2066	0.1964	0.959	1.0782	1.0316	0.3477	0.2921	0.950	0	
N&S	1.0140	1.0116	0.1847	0.1848	0.961	1.1099	1.1037	0.2987	0.3215	0.896	2	
SIMEX - Q	0.9891	0.9783	0.1838	0.1838	0.955	0.8718	0.8545	0.2241	0.2194	0.840	0	
SIMEX - L	0.9704	0.9600	0.1769	0.1794	0.945	0.7587	0.7491	0.1799	0.1735	0.657	0	
SIMEX - NL	1.0223	1.0131	0.2497	0.2053	0.944	1.0206	0.9698	0.3251	0.2758	0.942	102	
H&W	1.0370	1.0350	0.4937	0.2231	0.985	1.0410	0.9682	0.4373	0.3121	0.957	2	
MCMC $\eta = 10$	1.0271	1.0294	0.1981	0.2182	0.95	0.9768	0.9642	0.2461	0.2941	0.94	0	
MCMC $\eta = 100$	1.0687	1.0769	0.2165	0.2251	0.93	1.0950	1.0748	0.3028	0.3490	0.99	0	
				$\gamma_1 = 0.35$	$\gamma_2 = 0.25$	$\gamma_3 = 0.50$						
Naive	0.9292	0.9207	0.1676	0.1657	0.914	0.5091	0.5029	0.1282	0.1285	0.060	0	
RECS	1.0439	1.0182	0.2389	0.2105	0.962	1.1123	0.9989	0.4765	0.3793	0.929	0	
N&S	0.9982	0.9929	0.1835	0.1839	0.945	1.0293	1.0674	0.2451	0.2514	0.858	12	
SIMEX - Q	0.9662	0.9564	0.1797	0.1757	0.942	0.7259	0.7125	0.1986	0.1868	0.605	0	
SIMEX - L	0.9450	0.9362	0.1721	0.1720	0.928	0.6028	0.5954	0.1534	0.1520	0.230	0	
SIMEX - NL	1.0190	0.9887	0.3647	0.2187	0.922	0.9651	0.8793	0.4155	0.2943	0.874	67	
H&W	1.1080	1.0940	0.6033	0.2595	0.981	1.0600	0.9600	0.5004	0.3425	0.942	3	
MCMC $\eta = 10$	1.0072	1.0051	0.1939	0.2109	0.93	0.8949	0.8863	0.2432	0.2653	0.86	0	
MCMC $\eta = 100$	1.0789	1.0910	0.2335	0.2403	0.95	1.1408	1.0934	0.3828	0.3606	0.98	0	

N&S - Novick and Stefanski (2002); H&W - Huang and Wang (2001)

SIMEX results are based on $B = 100$, $\lambda = (0.5, 0.7, 0.9, 1.1, 1.3, 1.5, 1.7, 1.9)$ and the simex R library

L - linear extrapolation; Q - quadratic extrapolation; NL - non-linear extrapolation of Cook and Stefanski (1994)

M - empirical mean; MD - empirical median; Emp-SD - empirical standard deviation; IQ-SD - inter-quartile dispersion; 95% CI - empirical coverage rate of 95% Wald confidence interval (or HPD interval, for MCMC) ; F - number of samples with no solution

Web Table 2

Simulation Study B
 Non-Normal Measurement Error with Normal or Non-Normal True Covariate
 Sample Size $n=200$, Validation Sample Size $m=70$

	β_0					β_1					F
	M	MD	Emp-SD	IQ-SD	95% CI	M	MD	Emp-SD	IQ-SD	95% CI	
$X \sim N(0, 1); \epsilon \sim \text{Double exponential with } \text{Var}(\epsilon) = 0.5$											
Naive	0.9467	0.9350	0.1775	0.1804	0.925	0.6378	0.6287	0.1609	0.1517	0.334	0
RECS	1.0288	1.0048	0.2192	0.1989	0.952	1.0684	0.9986	0.3611	0.3264	0.948	0
N&S	1.0040	0.9978	0.1883	0.1758	0.969	1.0471	1.0126	0.2970	0.3293	0.888	0
SIMEX - Q	0.9884	0.9763	0.1931	0.1883	0.952	0.8905	0.8641	0.2438	0.2276	0.871	0
SIMEX - L	0.9654	0.9529	0.1836	0.1772	0.940	0.7519	0.7381	0.1924	0.1831	0.652	0
SIMEX - NL	1.0471	1.0189	0.2754	0.2135	0.945	1.1111	1.0256	0.4207	0.3336	0.967	68
H&W	0.9980	0.9919	0.4298	0.2157	0.982	1.0770	0.9848	0.4523	0.3202	0.951	0
MCMC $\eta=10$	1.0208	1.0349	0.1957	0.2084	0.94	1.0349	1.0186	0.2258	0.2753	0.990	0
MCMC $\eta=100$	1.1028	1.0933	0.2216	0.2410	0.97	1.2987	1.2613	0.3796	0.4360	0.970	0
$X \sim N(0, 1); \epsilon \sim \text{Double exponential with } \text{Var}(\epsilon) = 1$											
Naive	0.9174	0.9078	0.1731	0.1787	0.911	0.4659	0.4598	0.1370	0.1349	0.039	0
RECS	1.0545	1.0167	0.2735	0.2291	0.962	1.1362	1.0089	0.5714	0.4335	0.933	6
N&S	0.9700	0.9647	0.1851	0.1758	0.934	0.9625	1.0099	0.2794	0.2872	0.845	11
SIMEX - Q	0.9583	0.9474	0.1891	0.1861	0.933	0.7205	0.6970	0.2289	0.2165	0.629	0
SIMEX - L	0.9327	0.9228	0.1782	0.1772	0.912	0.5599	0.5514	0.1659	0.1601	0.197	0
SIMEX - NL	1.0032	0.9982	0.5742	0.2513	0.790	1.2130	1.0485	0.7743	0.4804	0.909	106
H&W	0.9353	0.9704	0.6913	0.2543	0.973	0.9886	0.8985	0.4636	0.3336	0.942	1
MCMC $\eta=10$	0.9779	0.9699	0.1876	0.2032	0.92	0.8304	0.8075	0.2289	0.2788	0.840	0
MCMC $\eta=100$	1.0711	1.0612	0.2221	0.2204	0.96	1.1476	1.0940	0.3785	0.4483	0.990	0
$X \sim N(0, 1); \epsilon \sim \text{Modified } \chi^2_{(1)} \text{ with } \text{Var}(\epsilon) = 0.5$											
Naive	0.9732	0.9745	0.1785	0.1692	0.932	0.6822	0.6793	0.1706	0.1672	0.467	0
RECS	1.0335	1.0295	0.2091	0.1929	0.951	1.0555	1.0348	0.2981	0.2775	0.959	0
N&S	1.0636	1.0442	0.2100	0.2085	0.946	1.1449	1.1451	0.3230	0.3808	0.857	4
SIMEX - Q	1.0340	1.0277	0.2044	0.1846	0.949	0.9678	0.9500	0.2624	0.2617	0.890	0
SIMEX - L	0.9980	0.9973	0.1878	0.1727	0.942	0.8080	0.8028	0.2052	0.2024	0.720	0
SIMEX - NL	1.1297	1.0882	0.3964	0.2483	0.912	1.2028	1.1617	0.3912	0.3573	0.975	45
H&W	0.7934	0.8692	0.5560	0.1964	0.963	1.0260	0.9809	0.3822	0.2861	0.956	2
MCMC $\eta=10$	1.0813	1.0884	0.2050	0.2144	0.93	1.0596	1.0267	0.2486	0.2244	0.97	0
MCMC $\eta=100$	1.1501	1.1402	0.2368	0.2504	0.93	1.2154	1.1673	0.3372	0.2999	0.92	0
$X \sim N(0, 1); \epsilon \sim \text{Modified } \chi^2_{(1)} \text{ with } \text{Var}(\epsilon) = 1$											
Naive	0.9510	0.9502	0.1765	0.1667	0.926	0.5196	0.5146	0.1547	0.1552	0.126	0
RECS	1.0519	1.0315	0.2444	0.2116	0.961	1.0869	1.0345	0.3970	0.3546	0.966	0
N&S	1.0914	1.0811	0.2151	0.2102	0.909	1.1232	1.1779	0.2711	0.2319	0.805	18
SIMEX - Q	1.0262	1.0140	0.2135	0.1890	0.933	0.8320	0.8234	0.2711	0.2765	0.747	0
SIMEX - L	0.9734	0.9703	0.1857	0.1690	0.931	0.6269	0.6187	0.1880	0.1853	0.316	0
SIMEX - NL	1.1226	1.0980	0.1702	0.4804	0.435	1.6444	1.4262	0.9090	0.7413	0.958	59
H&W	0.5574	0.7494	0.8302	0.2506	0.934	0.9640	0.8949	0.4753	0.3069	0.927	3
MCMC $\eta=10$	1.0585	1.0639	0.1975	0.2187	0.95	0.9402	0.9125	0.2241	0.2299	0.940	0
MCMC $\eta=100$	1.1556	1.1341	0.2378	0.2447	0.94	1.1502	1.1252	0.2996	0.2924	0.980	0
$X \sim \text{Modified } \chi^2_{(1)} \text{ with } \text{Var}(X) = 1; \epsilon \sim \text{Modified } \chi^2_{(1)} \text{ with } \text{Var}(\epsilon) = 0.5$											
Naive	0.9059	0.9015	0.1689	0.1632	0.903	0.5461	0.5348	0.1709	0.1662	0.250	0
RECS	1.0862	1.0356	0.2953	0.2626	0.967	1.1279	1.0479	0.4989	0.4415	0.938	0
N&S	1.2102	1.1694	0.3294	0.3620	0.813	1.2750	1.3552	0.4993	0.6506	0.672	51
SIMEX - Q	0.9703	0.9592	0.1968	0.1913	0.937	0.8175	0.7884	0.2870	0.2713	0.791	0
SIMEX - L	0.9253	0.9166	0.1750	0.1690	0.924	0.6492	0.6354	0.2056	0.1979	0.459	0
SIMEX - NL	1.0547	1.0083	0.9484	0.3625	0.495	1.2668	1.1121	0.6949	0.5137	0.945	43
H&W	0.6861	0.8618	0.8291	0.2053	0.944	0.9610	0.8409	0.5507	0.3729	0.912	6
MCMC $\eta=10$	1.0400	1.0104	0.2344	0.1886	0.90	0.7529	0.7365	0.3067	0.3059	0.900	0
MCMC $\eta=100$	1.1630	1.1179	0.3110	0.2506	0.91	1.0279	1.0105	0.4865	0.4735	0.970	0
$X \sim \text{Modified } \chi^2_{(1)} \text{ with } \text{Var}(X) = 1; \epsilon \sim \text{Modified } \chi^2_{(1)} \text{ with } \text{Var}(\epsilon) = 1$											
Naive	0.8784	0.8740	0.1661	0.1604	0.866	0.3917	0.3824	0.1448	0.1402	0.063	0
RECS	1.1134	1.0370	0.3589	0.2897	0.967	1.1272	1.0319	0.5784	0.5316	0.932	0
N&S	1.1288	1.0985	0.2997	0.2986	0.651	1.0147	1.1508	0.4306	0.5522	0.545	212
SIMEX - Q	0.9337	0.9243	0.1926	0.1794	0.917	0.6396	0.6176	0.2602	0.2498	0.490	0
SIMEX - L	0.8923	0.8876	0.1709	0.1616	0.890	0.4720	0.4604	0.1749	0.1683	0.099	0
SIMEX - NL	0.7096	0.8059	0.8610	0.4040	0.451	1.4470	1.1876	1.0740	0.8718	0.827	146
H&W	0.6004	0.7683	0.7942	0.2090	0.917	0.8084	0.6995	0.5894	0.4255	0.846	13
MCMC $\eta=10$	0.9865	0.9440	0.2132	0.2141	0.93	0.5704	0.5703	0.2674	0.2727	0.700	0
MCMC $\eta=100$	1.1116	1.0870	0.2817	0.2235	0.96	0.8416	0.8245	0.4370	0.4278	0.930	0

N&S - Novick and Stefanski (2002); H&W - Huang and Wang (2001)

SIMEX results are based on $B = 100$, $\lambda = (0.5, 0.7, 0.9, 1.1, 1.3, 1.5, 1.7, 1.9)$ and the simex R library

L - linear extrapolation; Q - quadratic extrapolation; NL - non-linear extrapolation of Cook and Stefanski (1994)

M - empirical mean; MD - empirical median; Emp-SD - empirical standard deviation; IQ-SD - inter-quartile dispersion; 95% CI - empirical coverage rate of 95% Wald confidence interval (or HPD interval, for MCMC); F - number of samples with no solution

Web Table 3
Simulation Study C
Two Error-Prone Covariates and One Error-Free Covariate
Scenario C1: $X_1, X_2, Z \sim N(0, 1)$ and $\epsilon_1, \epsilon_2 \sim N(0, 1)$
Sample Size $n=500$

	β_1					β_2					
	M	MD	Emp-SD	IQ-SD	95% CI	M	MD	Emp-SD	IQ-SD	95% CI	
Naive	0.4296	0.4242	0.0804	0.0781	0.000	0.4282	0.4235	0.0853	0.0857	0.000	
RECS	1.0817	1.0341	0.3116	0.2595	0.972	1.0784	1.0305	0.3333	0.2764	0.956	
N&S	0.8286	0.8288	0.1316	0.1319	0.826	0.8230	0.8259	0.1444	0.1454	0.775	
SIMEX - Q	0.8113	0.8038	0.1377	0.1333	0.571	0.8096	0.8006	0.1450	0.1378	0.580	
SIMEX - L	0.5235	0.5189	0.0917	0.0911	0.003	0.5219	0.5150	0.0969	0.0953	0.000	
SIMEX - NL	-2.6768	-2.5286	1.1003	1.0482	0.000	-2.5988	-2.4974	1.1896	1.0356	0.000	
H&W	0.6678	0.6675	0.5038	0.3360	0.800	0.6975	0.6744	0.5318	0.3407	0.786	
MCMC $\eta=10$	0.7675	0.7560	0.1670	0.1550	0.550	0.7695	0.7706	0.1550	0.1188	0.640	
MCMC $\eta=100$	1.0608	1.0335	0.2765	0.2508	0.870	1.0688	1.0593	0.2804	0.1833	0.890	

	β_0					β_3					F
	M	MD	Emp-SD	IQ-SD	95% CI	M	MD	Emp-SD	IQ-SD	95% CI	
Naive	0.8587	0.8575	0.1163	0.1129	0.734	0.8626	0.8599	0.1201	0.1223	0.769	0
RECS	1.0500	1.0294	0.2112	0.1791	0.972	1.0525	1.0286	0.2079	0.1907	0.979	0
N&S	0.9432	0.9433	0.1379	0.1383	0.947	0.9483	0.9448	0.1382	0.1412	0.951	52
SIMEX - Q	0.9553	0.9502	0.1433	0.1394	0.886	0.9587	0.9524	0.1452	0.1519	0.920	0
SIMEX - L	0.8821	0.8803	0.1215	0.1214	0.783	0.8857	0.8819	0.1247	0.1278	0.809	0
SIMEX - NL	0.3396	0.3544	1.1798	0.5785	0.051	0.3642	0.4221	1.1742	0.5205	0.059	285
H&W	0.7535	0.8516	0.7885	0.2376	0.887	1.0277	0.9679	0.4669	0.3039	0.918	14
MCMC $\eta=10$	0.9101	0.9060	0.1349	0.1290	0.840	0.9172	0.9113	0.1281	0.1475	0.870	0
MCMC $\eta=100$	1.0407	1.0328	0.1964	0.1728	0.930	1.0445	1.0418	0.1622	0.1834	0.970	0

N&S - Novick and Stefanski (2002); H&W - Huang and Wang (2001)

SIMEX results are based on $B = 100$, $\lambda = (0.5, 0.7, 0.9, 1.1, 1.3, 1.5, 1.7, 1.9)$ and the simex R library

L - linear extrapolation; Q - quadratic extrapolation; NL - non-linear extrapolation of Cook and Stefanski (1994)

M - empirical mean; MD - empirical median; Emp-SD - empirical standard deviation; IQ-SD - inter-quartile dispersion; 95% CI - empirical coverage rate of 95% Wald confidence interval (or HPD interval, for MCMC); F - number of samples with no solution

Web Table 4

Simulation Study C

Two Error-Prone Covariates and One Error-Free Covariate

Scenario C2: $X_1, X_2, Z \sim N(0, 1)$ and $\epsilon_1, \epsilon_2 \sim N(0, 0.4 + 0.25(|X_1| + |X_2| + |Z|))$

Sample Size $n=500$

	β_1					β_2					
	M	MD	Emp-SD	IQ-SD	95% CI	M	MD	Emp-SD	IQ-SD	95% CI	
Naive	0.4386	0.4334	0.0824	0.0792	0.001	0.4364	0.4313	0.0868	0.0857	0.000	
RECS	1.1884	1.1122	0.3800	0.3126	0.986	1.1822	1.1151	0.4044	0.3313	0.977	
N&S	0.8606	0.8619	0.1329	0.1312	0.842	0.8521	0.8559	0.1468	0.1454	0.834	
SIMEX - Q	0.8345	0.8242	0.1431	0.1394	0.819	0.8315	0.8231	0.1497	0.1439	0.767	
SIMEX - L	0.5339	0.5265	0.0940	0.0903	0.009	0.5314	0.5257	0.0985	0.0973	0.011	
SIMEX - NL	-2.3990	-2.2229	0.9623	0.9407	0.000	-2.3830	-2.1980	1.0071	0.9666	0.000	
H&W	0.6165	0.6176	0.4809	0.3277	0.733	0.6455	0.6072	0.4809	0.3271	0.766	
MCMC $\eta=10$	0.7713	0.7621	0.1663	0.1656	0.600	0.7716	0.7630	0.1606	0.1188	0.620	
MCMC $\eta=100$	1.0338	1.0084	0.2709	0.2637	0.890	1.0378	1.0093	0.2576	0.1849	0.900	

	β_0					β_3					F
	M	MD	Emp-SD	IQ-SD	95% CI	M	MD	Emp-SD	IQ-SD	95% CI	
Naive	0.8593	0.8586	0.1162	0.1153	0.740	0.8712	0.8694	0.1223	0.1223	0.791	0
RECS	1.0787	1.0475	0.2359	0.1896	0.981	1.0828	1.0482	0.2405	0.2007	0.983	1
N&S	0.9353	0.9350	0.1360	0.1387	0.939	0.9668	0.9655	0.1439	0.1465	0.966	46
SIMEX - Q	0.9569	0.9531	0.1434	0.1394	0.907	0.9793	0.9746	0.1516	0.1562	0.933	0
SIMEX - L	0.8827	0.8824	0.1214	0.1219	0.808	0.8961	0.8921	0.1276	0.1314	0.834	0
SIMEX - NL	0.3789	0.3653	1.1233	0.5524	0.035	0.3096	0.3920	0.9646	0.4150	0.039	185
H&W	0.7637	0.8325	0.9796	0.2280	0.887	0.9627	0.9279	0.4248	0.2837	0.905	9
MCMC $\eta=10$	0.9148	0.9051	0.1358	0.1285	0.830	0.9237	0.9205	0.1262	0.1474	0.890	0
MCMC $\eta=100$	1.0341	1.0186	0.1903	0.1847	0.920	1.0358	1.0339	0.1626	0.1701	0.950	0

N&S - Novick and Stefanski (2002); H&W - Huang and Wang (2001)

SIMEX results are based on $B = 100$, $\lambda = (0.5, 0.7, 0.9, 1.1, 1.3, 1.5, 1.7, 1.9)$ and the simex R library

L - linear extrapolation; Q - quadratic extrapolation; NL - non-linear extrapolation of Cook and Stefanski (1994)

M - empirical mean; MD - empirical median; Emp-SD - empirical standard deviation; IQ-SD - inter-quartile dispersion; 95% CI - empirical coverage rate of 95% Wald confidence interval (or HPD interval, for MCMC); F - number of samples with no solution

Web Table 5

Simulation Study C
 Two Error-Prone Covariates and One Error-Free Covariate
 Scenario C3: $X_1, X_2 \sim \text{Modified } \chi_1^2$, $\text{Var}(X_1) = \text{Var}(X_2) = 1$, $Z \sim N(0, 1)$,
 and $\epsilon_1, \epsilon_2 \sim \text{Modified } \chi_1^2$, $\text{Var}(\epsilon_1) = \text{Var}(\epsilon_2) = 1$
 Sample Size $n=500$

	β_1					β_2					
	M	MD	Emp-SD	IQ-SD	95% CI	M	MD	Emp-SD	IQ-SD	95% CI	
Naive	0.3853	0.3814	0.0887	0.0915	0.000	0.3830	0.3811	0.0843	0.0856	0.000	
RECS	1.0430	0.9841	0.3726	0.3328	0.942	1.0322	0.9850	0.3319	0.3016	0.946	
N&S	0.8656	0.8664	0.2576	0.2880	0.782	0.8659	0.8750	0.2516	0.2872	0.784	
SIMEX - Q	0.7463	0.7315	0.1586	0.1598	0.445	0.7431	0.7317	0.1493	0.1488	0.450	
SIMEX - L	0.4672	0.4624	0.1009	0.1021	0.001	0.4647	0.4602	0.0958	0.0989	0.000	
SIMEX - NL	-1.5004	-1.2992	0.7297	0.5271	0.000	-1.5046	-1.3341	0.6896	0.5078	0.000	
H&W	0.4723	0.4276	0.2733	0.2108	0.654	0.4625	0.4216	0.2849	0.1992	0.606	
MCMC $\eta=10$	0.7216	0.7191	0.1696	0.2000	0.640	0.7054	0.7066	0.1493	0.1446	0.600	
MCMC $\eta=100$	1.1462	1.1199	0.3153	0.3259	0.910	1.1100	1.0941	0.3072	0.2442	0.950	
	β_0					β_3					
	M	MD	Emp-SD	IQ-SD	95% CI	M	MD	Emp-SD	IQ-SD	95% CI	F
Naive	0.8007	0.7971	0.1144	0.1113	0.548	0.8883	0.8893	0.1206	0.1207	0.809	0
RECS	1.0595	1.0225	0.2447	0.2084	0.963	1.0413	1.0232	0.1856	0.1667	0.971	0
N&S	1.0389	1.0389	0.1554	0.1553	0.945	0.9261	0.9201	0.1396	0.1415	0.946	34
SIMEX - Q	0.9049	0.8978	0.1456	0.1383	0.823	0.9626	0.9588	0.1402	0.1399	0.915	0
SIMEX - L	0.8199	0.8146	0.1191	0.1133	0.598	0.9059	0.9073	0.1243	0.1244	0.893	0
SIMEX - NL	0.5453	0.5554	0.1777	0.1286	0.027	0.4842	0.5551	0.9577	0.3875	0.122	60
H&W	0.3332	0.4979	0.6451	0.2517	0.681	0.9155	0.9023	0.1798	0.1638	0.971	1
MCMC $\eta=10$	0.9526	0.9413	0.1204	0.0946	0.900	0.9125	0.8945	0.1388	0.1345	0.840	0
MCMC $\eta=100$	1.2427	1.2461	0.1978	0.2014	0.830	1.0167	0.9887	0.1713	0.1519	0.910	0

N&S - Novick and Stefanski (2002); H&W - Huang and Wang (2001)
 SIMEX results are based on $B = 100$, $\lambda = (0.5, 0.7, 0.9, 1.1, 1.3, 1.5, 1.7, 1.9)$ and the simex R library
 L - linear extrapolation; Q - quadratic extrapolation; NL - non-linear extrapolation of Cook and Stefanski (1994)
 M - empirical mean; MD - empirical median; Emp-SD - empirical standard deviation; IQ-SD - inter-quartile dispersion;
 95% CI - empirical coverage rate of 95% Wald confidence interval (or HPD interval, for MCMC); F - number of samples
 with no solution

Web Table 6

Simulation Study D
 Performance of RECS Under Misspecified Error Model
 Assumed Error Model: $\epsilon \sim N(0, \sigma^2)$
 Results for Estimate of β_1

Distn of X	True Distn of ϵ	Naive Estimate				RECS Estimate			
		Mean	Median	Emp-SD	IQ-SD	Mean	Median	Emp-SD	IQ-SD
Normal	Skewed Normal	0.6621	0.6569	0.1655	0.1668	1.1885	1.1028	0.4790	0.3466
Normal	MODCHI	0.6821	0.6791	0.1706	0.1672	1.2778	1.1592	0.5621	0.4267
Skewed Normal	Skewed Normal	0.5992	0.5895	0.1620	0.1589	1.1983	1.0923	0.5506	0.4355
Skewed Nomal	MODCHI	0.6058	0.6043	0.1669	0.1698	1.3466	1.1862	0.6752	0.5526
MODCHI	Skewed Normal	0.5250	0.5209	0.1573	0.1532	1.3280	1.1335	0.7406	0.5932
MODCHI	MODCHI	0.5460	0.5347	0.1710	0.1661	1.6553	1.3567	1.0210	0.9698